

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human KIR3DL3/CD158z in direct ELISAs.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 1136B
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Human embryonic kidney cell line HEK293-derived recombinant human KIR3DL3/CD158z Met1-Leu322 Accession # Q8N743
<b>Formulation</b>	Supplied as a solution in PBS containing BSA, Glycerol and Sodium Azide. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	1:1000 dilution	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C, as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after opening.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after opening.</li> </ul>

**BACKGROUND**

KIR3DL3 is a member of the killer-immunoglobulin-like receptors (KIR), a functionally diverse family of transmembrane receptors expressed primarily on NK cells. KIR3DL3 is a "framework" KIR gene, and is present in all individuals, in contrast to non-framework KIRs, whose gene content varies by KIR haplotype. KIR3DL3 is expressed at low or undetectable levels in peripheral blood NK cells, but expression can be induced in NK cell lines by demethylation agents, indicating that promoter DNA methylation inhibits KIR3DL3 transcription. Despite the lack of evidence for surface protein expression, KIR3DL3 mRNA expression has been detected in both decidual and peripheral blood NK cells in normal, disease-free conditions.

**PRODUCT SPECIFIC NOTICES**

\* Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to SDS for additional information and handling instructions.