

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human eNOS when phosphorylated at S1177 in direct ELISAs and Western blots.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 1014F
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	Phosphopeptide containing the human eNOS S1177 site Accession # P29474
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.5 µg/mL	See Below

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

## BACKGROUND

Endothelial NOS (eNOS), also known as nitric oxide synthase 3 (NOS3) or constitutive NOS (cNOS), is an enzyme encoded by the NOS3 gene. Endothelial NOS generates nitric oxide in blood vessels and is involved with regulating vascular tone by inhibiting smooth muscle contraction and platelet aggregation. A constitutive calcium dependent NOS provides a basal release of NO. eNOS is associated with plasma membranes surrounding cells and the membranes of the Golgi apparatus within cells.