

DESCRIPTION



Species Reactivity	Human
Specificity	Detects human SUSD2 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 944812
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell line HEK293-derived human SUSD2 Met1-Ala785 Accession # Q9UGT4
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	See Below
Immunohistochemistry	8-25 µg/mL	See Below

DATA

<p>Western Blot</p>  <p>Detection of Human SUSD2 by Western Blot. Western blot shows lysates of SK-BR-3 human breast cancer cell line. PVDF membrane was probed with 0.1 µg/mL of Mouse Anti-Human SUSD2 Monoclonal Antibody (Catalog # MAB9056) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # HAF018). Specific bands were detected for SUSD2 at approximately 60 and 110 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p>Immunohistochemistry</p>  <p>SUSD2 in Human Breast Cancer Tissue. SUSD2 was detected in immersion fixed paraffin-embedded sections of human breast cancer tissue using Mouse Anti-Human SUSD2 Monoclonal Antibody (Catalog # MAB9056) at 15 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Mouse HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS002) and counterstained with hematoxylin (blue). Specific staining was localized to cytoplasm with higher expression in cancer tissue. View our protocol for Chromogenic IHC Staining of Paraffin-embedded Tissue Sections.</p>
---	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Sushi domain containing 2, or SUSD2, is a type I transmembrane protein of 822 amino acids containing functional domains inherent to adhesion molecules. SUSD2 has been described as a novel marker of human endometrial mesenchymal stem-like cells and it has been used for their prospective isolation. As a transmembrane receptor, SUSD2 has been proposed to interact with Galectin-1 and to be the receptor for C10ORF99, a novel potential cytokine suggested to inhibit colon cancer cell growth through inducing G1 arrest. There is evidence that SUSD2 may play a role in breast tumorigenesis.