

#### DESCRIPTION


<b>Species Reactivity</b>	Human/Rat
<b>Specificity</b>	Detects human SUSD2 in direct ELISAs and Western blots.
<b>Source</b>	Recombinant Monoclonal Rabbit IgG Clone # 1279D
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	HEK293 human embryonic kidney cell line-derived human SUSD2 Met1-Ala785 Accession # Q9UGT4
<b>Formulation</b>	Supplied as a solution in PBS containing BSA, Glycerol and Sodium Azide. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below
<b>Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	See Below

#### DATA



#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C, as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after opening.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after opening.</li> </ul>

#### BACKGROUND

Sushi domain containing 2, or SUSD2, is a type I transmembrane protein of 822 amino acids containing functional domains inherent to adhesion molecules. SUSD2 has been described as a novel marker of human endometrial mesenchymal stem-like cells and it has been used for their prospective isolation. As a transmembrane receptor, SUSD2 has been proposed to interact with Galectin-1 and to be the receptor for C10ORF99, a novel potential cytokine suggested to inhibit colon cancer cell growth through inducing G1 arrest. There is evidence that SUSD2 may play a role in breast tumorigenesis.

#### PRODUCT SPECIFIC NOTICES

Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to SDS for additional information and handling instructions.