

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human IL-18/IL-1F4 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2A} Clone # 914205
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human IL-18/IL-1F4 synthetic peptide Accession # Q14116
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Neutralization	Measured by its ability to neutralize IL-18/IL-1F4-induced IFN-γ secretion in the KG-1 human acute myelogenous leukemia cell line. Novick, D. <i>et al.</i> (1999) <i>Immunity</i> 10 (1):127. The Neutralization Dose (ND ₅₀) is typically 3-15 µg/mL in the presence of 10 ng/mL Recombinant Human IL-18/IL-1F4 and 20 ng/mL Recombinant Human TNF-α.
-----------------------	--

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Interleukin-18 (IL-18) is a proinflammatory cytokine in the IL-1 family that exerts distinct immune effects depending on the local cytokine environment. It is expressed as a 24 kDa precursor by endothelial and epithelial cells, keratinocytes, $\gamma\delta$ T cells, and phagocytes. The precursor is activated intracellularly by Caspase-1 mediated proteolysis to release the 17 kDa mature cytokine. The precursor can also be released by necrotic cells for extracellular cleavage by multiple proteases. IL-18 activation is induced by infection or tissue damage and contributes to disease pathology in chronic inflammation (1-3). IL-18 binds to the widely expressed IL-18 R α which recruits IL-18 R β to form the signaling receptor complex (4, 5). Its bioactivity is negatively regulated by interactions with IL-18 binding proteins and virally encoded IL-18BP homologs (6). In the presence of IL-12 or IL-15, IL-18 enhances anti-viral Th1 immune responses by inducing IFN- γ production and the cytolytic activity of CD8⁺ T cells and NK cells (7, 8). In the absence of IL-12 or IL-15, however, IL-18 promotes production of the Th2 cytokines IL-4 and IL-13 by CD4⁺ T cells and basophils (9, 10). In the presence of IL-1 β or IL-23, IL-18 induces the antigen-independent production of IL-17 by $\gamma\delta$ T cells and CD4⁺ T cells (11). IL-18 also promotes myeloid dendritic cell maturation and triggers neutrophil respiratory burst (12, 13). In cancer, IL-18 exhibits diverse activities including enhancing anti-tumor immunity, inhibiting or promoting angiogenesis, and promoting tumor cell metastasis (14). Mature human IL-18 shares approximately 63% amino acid sequence identity with mouse and rat IL-18 (15). Alternative splicing in human ovarian cancer generates an isoform that is resistant to Caspase-1 activation (16). A cell surface form can be expressed on M-CSF induced macrophages and released in response to bacterial endotoxin (17).

References:

1. Dinarello, C.A. et al. (2013) *Front. Immunol.* **4**:289.
2. Smith, D.E. (2011) *J. Leukoc. Biol.* **89**:383.
3. Gu, Y. et al. (1997) *Science* **275**:206.
4. Torigoe, K. et al. (1997) *J. Biol. Chem.* **272**:25737.
5. Cheung, H. et al. (2005) *J. Immunol.* **174**:5351.
6. Novick, D. et al. (1999) *Immunity* **10**:127.
7. Fehniger, T.A. et al. (1999) *J. Immunol.* **162**:4511.
8. Yoshimoto, T. et al. (1998) *J. Immunol.* **161**:3400.
9. Yoshimoto, T. et al. (2000) *Nat. Immunol.* **1**:132.
10. Kroeger, K.M. et al. (2009) *J. Leukoc. Biol.* **86**:769.
11. Lalor, S.J. et al. (2011) *J. Immunol.* **186**:5738.
12. Li, J. et al. (2004) *Cell. Immunol.* **227**:103.
13. Elbim, C. et al. (2005) *Clin. Diagn. Lab. Immunol.* **12**:436.
14. Fabbri, M. et al. (2015) *J. Leukoc. Biol.* **97**:665.
15. Ushio, S. et al. (1996) *J. Immunol.* **156**:4274.
16. Gaggero, A. et al. (2004) *Oncogene* **23**:7552.
17. Bellora, F. et al. (2012) *Eur. J. Immunol.* **42**:1618.