

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-18/IL-1F4 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 923503
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human IL-18/IL-1F4 Tyr37-Asp193 Accession # Q14116
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	5-25 µg/mL	See Below


DATA

Immunohistochemistry



IL-18/IL-1F4 in Human Spleen. IL-18/IL-1F4 was detected in immersion fixed paraffin-embedded sections of human spleen using Mouse Anti-Human IL-18/IL-1F4 Monoclonal Antibody (Catalog # MAB91245) at 15 µg/mL for 1 hour at room temperature followed by incubation with the Anti-Mouse IgG VisUCyte™ HRP Polymer Antibody (Catalog # Catalog # VC001). Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # Catalog # CTS013). Tissue was stained using DAB (brown) and counterstained with hematoxylin (blue). Specific staining was localized to T cells. View our protocol for IHC Staining with VisUCyte HRP Polymer Detection Reagents.

Immunohistochemistry



IL-18/IL-1F4 in Human Pancreas. IL-18/IL-1F4 was detected in immersion fixed paraffin-embedded sections of human pancreas using Mouse Anti-Human IL-18/IL-1F4 Monoclonal Antibody (Catalog # MAB91245) at 5 µg/mL for 1 hour at room temperature followed by incubation with the Anti-Mouse IgG VisUCyte™ HRP Polymer Antibody (Catalog # Catalog # VC001). Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # Catalog # CTS013). Tissue was stained using DAB (brown) and counterstained with hematoxylin (blue). Specific staining was localized to cytoplasm in islet cells. View our protocol for IHC Staining with VisUCyte HRP Polymer Detection Reagents.

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Interleukin-18 (IL-18) is a proinflammatory cytokine in the IL-1 family that exerts distinct immune effects depending on the local cytokine environment. It is expressed as a 24 kDa precursor by endothelial and epithelial cells, keratinocytes, $\gamma\delta$ T cells, and phagocytes. The precursor is activated intracellularly by Caspase-1 mediated proteolysis to release the 17 kDa mature cytokine. The precursor can also be released by necrotic cells for extracellular cleavage by multiple proteases. IL-18 activation is induced by infection or tissue damage and contributes to disease pathology in chronic inflammation (1-3). IL-18 binds to the widely expressed IL-18 R α which recruits IL-18 R β to form the signaling receptor complex (4, 5). Its bioactivity is negatively regulated by interactions with IL-18 binding proteins and virally encoded IL-18BP homologs (6). In the presence of IL-12 or IL-15, IL-18 enhances anti-viral Th1 immune responses by inducing IFN- γ production and the cytolytic activity of CD8⁺ T cells and NK cells (7, 8). In the absence of IL-12 or IL-15, however, IL-18 promotes production of the Th2 cytokines IL-4 and IL-13 by CD4⁺ T cells and basophils (9, 10). In the presence of IL-1 β or IL-23, IL-18 induces the antigen-independent production of IL-17 by $\gamma\delta$ T cells and CD4⁺ T cells (11). IL-18 also promotes myeloid dendritic cell maturation and triggers neutrophil respiratory burst (12, 13). In cancer, IL-18 exhibits diverse activities including enhancing anti-tumor immunity, inhibiting or promoting angiogenesis, and promoting tumor cell metastasis (14). Mature human IL-18 shares approximately 63% amino acid sequence identity with mouse and rat IL-18 (15). Alternative splicing in human ovarian cancer generates an isoform that is resistant to Caspase-1 activation (16). A cell surface form can be expressed on M-CSF induced macrophages and released in response to bacterial endotoxin (17).

References:

1. Dinarello, C.A. *et al.* (2013) *Front. Immunol.* **4**:289.
2. Smith, D.E. (2011) *J. Leukoc. Biol.* **89**:383.
3. Gu, Y. *et al.* (1997) *Science* **275**:206.
4. Torigoe, K. *et al.* (1997) *J. Biol. Chem.* **272**:25737.
5. Cheung, H. *et al.* (2005) *J. Immunol.* **174**:5351.
6. Novick, D. *et al.* (1999) *Immunity* **10**:127.
7. Fehniger, T.A. *et al.* (1999) *J. Immunol.* **162**:4511.
8. Yoshimoto, T. *et al.* (1998) *J. Immunol.* **161**:3400.
9. Yoshimoto, T. *et al.* (2000) *Nat. Immunol.* **1**:132.
10. Kroeger, K.M. *et al.* (2009) *J. Leukoc. Biol.* **86**:769.
11. Lalor, S.J. *et al.* (2011) *J. Immunol.* **186**:5738.
12. Li, J. *et al.* (2004) *Cell. Immunol.* **227**:103.
13. Elbim, C. *et al.* (2005) *Clin. Diagn. Lab. Immunol.* **12**:436.
14. Fabb, M. *et al.* (2015) *J. Leukoc. Biol.* **97**:665.
15. Ushio, S. *et al.* (1996) *J. Immunol.* **156**:4274.
16. Gaggero, A. *et al.* (2004) *Oncogene* **23**:7552.
17. Bellora, F. *et al.* (2012) *Eur. J. Immunol.* **42**:1618.