

**DESCRIPTION**



<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse SPARC/Osteonectin in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human SPARC is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2B</sub> Clone # 124413
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse SPARC/Osteonectin Ala18-Ile302 Accession # P07214
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	2 µg/mL	See Below
<b>Immunohistochemistry</b>	8-25 µg/mL	See Below
<b>Intracellular Staining by Flow Cytometry</b>	0.25 µg/10 <sup>6</sup> cells	Balb/3T3 mouse embryonic fibroblast cell line fixed with paraformaldehyde and permeabilized with saponin
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**

<p><b>Western Blot</b></p>  <p><b>Detection of Mouse SPARC by Western Blot.</b> Western blot shows lysates of C2C12 mouse myoblast cell line and mouse placenta tissue. PVDF membrane was probed with 2 µg/mL of Rat Anti-Mouse SPARC Monoclonal Antibody (Catalog # MAB942) followed by HRP-conjugated Anti-Rat IgG Secondary Antibody (Catalog # HAF005). A specific band was detected for SPARC at approximately 35-37 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.</p>	<p><b>Immunohistochemistry</b></p>  <p><b>SPARC/Osteonectin in Mouse Ovary.</b> SPARC/Osteonectin was detected in perfusion fixed frozen sections of mouse ovary using Mouse SPARC/Osteonectin Monoclonal Antibody (Catalog # MAB942) at 25 µg/mL overnight at 4 °C. Tissue was stained using the Anti-Rat HRP-DAB Cell &amp; Tissue Staining Kit (brown; Catalog # CTS017) and counterstained with hematoxylin (blue). View our protocol for Chromogenic IHC Staining of Frozen Tissue Sections.</p>
--	--

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

SPARC, an acronym for "secreted protein, acidic and rich in cysteine", is also known as osteonectin or BM-40 (1-5). It is the founding member of a family of secreted matricellular proteins with similar domain structure. The 302 amino acid (aa), 43 kDa protein contains a 17 aa signal sequence, an N-terminal acidic region that binds calcium, a follistatin domain containing Kazal-like sequences, and a C-terminal extracellular calcium (EC) binding domain with two EF-hand motifs (1-5). Crystal structure shows that residues implicated in cell binding, inhibition of cell spreading and disassembly of focal adhesions cluster on one face of SPARC, while a collagen binding epitope and an N-glycosylation site are opposite this face (6). SPARC is produced by fibroblasts, capillary endothelial cells, platelets and macrophages, especially in areas of tissue morphogenesis and remodeling (3, 7). SPARC shows context-specific effects, but generally inhibits adhesion, spreading and proliferation, and promotes collagen matrix formation (3-5). For endothelial cells, SPARC disrupts focal adhesions and binds and sequesters PDGF and VEGF (3-5). SPARC is abundantly expressed in bone, where it promotes osteoblast differentiation and inhibits adipogenesis (5, 8). SPARC is potentially cleaved by metalloproteinases, producing an angiogenic peptide that includes the copper-binding sequence KGHK (7). Paradoxically, SPARC is highly expressed in many tumor types, yet expression mainly decreases the likelihood of metastasis and confers sensitivity to chemotherapy and radiation (4, 9, 10). Stabilin-1, which is expressed on alternately activated macrophages, is the first SPARC receptor to be identified. It binds the SPARC EC domain and mediates endocytosis for degradation (11). Mature mouse SPARC shows 97%, 92%, 92%, 92% and 83% aa identity with rat, human, dog, cow and chick SPARC, respectively.

**References:**

1. Lankat-Buttgereit, B. *et al.* (1988) FEBS Lett. **236**:352.
2. McVey, J.H. *et al.* (1988) J. Biol. Chem. **263**:11111.
3. Sage, H. *et al.* (1989) J. Cell Biol. **109**:341.
4. Framson, P.E. and E.H. Sage (2004) J. Cell. Biochem. **92**:679.
5. Alford, A.I. and K.D. Hankenson (2006) Bone **38**:749.
6. Hohenester, E. *et al.* (1997) EMBO J. **16**:3778.
7. Sage, E.H. *et al.* (2003) J. Biol. Chem. **278**:37849.
8. Delany, A.M. *et al.* (2003) Endocrinology **144**:2588.
9. Koblinski, J.E. *et al.* (2005) Cancer Res. **65**:7370.
10. Tai, I.T. *et al.* (2005) J. Clin. Invest. **115**:1492.
11. Kzhyshkowska, J. *et al.* (2006) J. Immunol. **176**:5825.