

DESCRIPTION

Species Reactivity	Rabbit
Specificity	Rabbit IgG (H+L)
Source	Polyclonal Donkey IgG
Purification	Antigen Affinity-purified
Immunogen	Rabbit IgG
Formulation	Supplied in a saline solution containing BSA and Sodium Azide.

*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Immunocytochemistry	Use at a 1:200 dilution
----------------------------	-------------------------

DATA

Immunocytochemistry



A172 (Positive) cells
No Primary Control

Detection of Rabbit IgG Primary Antibody by Immunocytochemistry.
 Detection of human Ki-67 by incubating cells with 3 µg/mL of Rabbit anti-Human Ki-67 monoclonal antibody (MAB7617) at room temperature for 1 hour followed by incubation with 1:200 dilution of Donkey Anti-Rabbit IgG NorthernLights™ NL557-conjugated secondary antibody for 1 hour at room temperature. Note strong positive staining in A172 cells (left panel, positive control) and a lack of non-specific background staining in the absence of primary antibodies (right panel, negative control).

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. <ul style="list-style-type: none"> • 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

R&D Systems offers a range of secondary antibodies and controls for flow cytometry, immunohistochemistry, and Western blotting. We provide species-specific secondary antibodies that are available with a variety of conjugated labels. Our NorthernLights fluorescent secondary antibodies are bright and resistant to photobleaching. We are currently offering secondary antibodies recognizing mouse, rat, goat, sheep, and rabbit IgG as well as chicken IgY. These reagents are available with three distinct excitation and emission maxima, making them ideal for multi-color fluorescence microscopy.