

Monoclonal Anti-human GR common/NR3C1 Antibody

ORDERING INFORMATION

Catalog Number: PP-H8004-00

Clone: H8004

GenBank: X03225

Ig Class: mouse IgG_{2A}

Volume: 100 µL

Concentration: 1 mg/mL

Formulation: A liquid formulation in physiologic saline with 0.1% NaN₃

Storage: ≤ -20 °C

Specificity: human GR common

Applications: Western Blot
Direct ELISA
Immunohistochemistry
Immunoprecipitation

Description

Glucocorticoid Receptor (GR; NR3C1) is a member of the Steroid Receptor family. It is expressed in almost all human tissues and organs. GR binds to its sequences as a homodimer. Two different forms, GR α and GR β , differing in their C-terminal parts were isolated. GR α is the classic GR that binds to glucocorticoids and transactivates or transrepresses glucocorticoid-responsive promoters. GR β does not bind glucocorticoids.

Preparation

Produced in BALB/c mouse ascites after inoculation with hybridoma of mouse myeloma cells (NS-1) and spleen cells derived from a BALB/c mouse immunized with recombinant human GR α (amino acids 2 - 60). The IgG fraction of the ascites was purified by ammonium sulfate fractionation.

Formulation

A liquid formulation in physiologic saline with 0.1% NaN₃.

Storage

This antibody is stable for greater than six months when held at -20 °C in a **manual defrost freezer** or at -70 °C. Upon thawing, the antibody can be stored at 2-8 °C for at least 1 month without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody specifically recognizes human GR α and GR β and cross-reacts with rat GR α and GR β . Not yet tested in other species.

Applications

Western Blot - This antibody can be used at 1-10 µg/mL under reducing conditions with the appropriate secondary reagents to detect human GR.

Direct ELISA - This antibody can be used at 0.35 µg/mL with the appropriate secondary reagents to detect human GR.

Immunohistochemistry - This antibody can be used at 10 µg/mL with the appropriate secondary reagents to detect human GR.

Immunoprecipitation - Optimal dilutions should be determined by each laboratory.

Optimal dilutions should be determined by each laboratory for each application.

Caution: Sodium azide may react with lead and copper plumbing to form explosive metal azides. Flush with large amounts of water during disposal.



Manufactured by:

Perseus Proteomics, Inc.
3F, A.i. Nihombashi EAST
30-1 Nihonbashi-Hakozakicho, Chuo-ku,
Tokyo 103-0015, JAPAN
Tel: +81-3-6264-8268
Fax: +81-3-3668-7776
E-mail: info@ppmx.com
http://www.ppmx.com

725968.1

9/16

FOR RESEARCH USE ONLY.
NOT FOR USE IN HUMANS.