

**MATERIAL DATA SHEET****Poly-Ub WT Chains (2-7) (K48-linked)****Cat. # UC-230**

Polyubiquitin chains are composed of ubiquitin monomers that are covalently linked through isopeptide bonds, which typically form between a lysine residue of one ubiquitin molecule and the C-terminal glycine residue of another ubiquitin molecule. Each human ubiquitin monomer is 76 amino acids (aa) in length and shares 96% and 100% aa identity with yeast and mouse ubiquitin, respectively. Seven of the 76 aa in ubiquitin are lysine residues that can participate in polyubiquitin chain formation. Linkage through specific lysine residues is thought to serve as a signal that affects protein degradation, signaling, trafficking, and other cellular processes. Linkage specific polyubiquitin chains are used to investigate mechanisms of chain recognition, binding and hydrolysis by the proteasome, deubiquitinating enzymes, E3 ligases or other proteins that contain ubiquitin-associated domains (UBAs) or ubiquitin-interacting motifs (UIMs). K48-linked chains are abundant in vivo and act as a universal signal for proteasomal degradation. This product is formed with wild-type human recombinant ubiquitin and linkage-specific enzymes. This mixture of polyubiquitin chains contains diubiquitin and higher MW species, monoubiquitin has been removed.

**Product Information**

<b>Quantity:</b>	100 µg, lyophilized powder
<b>Solubility:</b>	Reconstitute in aqueous buffer at 2 mg/ml
<b>Purity:</b>	> 95% by SDS-PAGE under reducing conditions and visualized by Colloidal Coomassie Blue stain.
<b>MW:</b>	17 kDa (Ub2) 26 kDa (Ub3), 34 kDa (Ub4), 43 kDa (Ub5), 52 kDa (Ub6), and 60 kDa (Ub7)

**Use & Storage**

<b>Use:</b>	Ubiquitin chains vary in length, linkage, and function. K48-linked polyubiquitin chains (Ub2-7) are ideal for investigating ubiquitin-binding proteins and as substrates for ubiquitin-specific isopeptidases. Reaction conditions will need to be optimized for each specific application. <b>IMPORTANT:</b> Heating this product in SDS-PAGE buffer or terminating reactions containing this product with heated SDS-PAGE buffer could lead to unexpected, high apparent molecular weight banding or smearing on gels that is not representative of product purity. For optimal results, we recommend incubation in SDS-PAGE buffer + DTT at <40 °C for 20 minutes prior to gel electrophoresis
<b>Storage:</b>	Solubilized solution at -20°C. Avoid multiple freeze/thaw cycles.

## Literature

- References:**
- Behrends, C. & J.W. Harper (2011) Nat. Struct. Mol. Biol. **18**: 520
  - Greene, W. *et al.* (2012) PLoS Pathog. **8**: e1002703
  - Henry, A.G. *et al.* (2012) Dev. Cell **23**: 519
  - Scheffner, M. *et al.* (1995) Nature **373**: 81
  - Sharp, P.M. & W.-H. Li (1987) Trends Ecol. Evol. **2**: 328
  - Tong, X. *et al.* (2012) J. Biol. Chem. **287**: 25280
  - Wei, W. *et al.* (2004) Nature **428**: 194
  - Zhang, J. *et al.* (2012) J. Biol. Chem. **287**: 28646

*For Laboratory Research Use Only, Not For Use in Humans*

Rev: 03/17/2014

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565

[www.bostonbiochem.com](http://www.bostonbiochem.com)

The contents of this datasheet (unless otherwise noted) are Copyright © 2008 Boston Biochem, Inc. All rights reserved. Duplication in whole or in part is strictly prohibited without the express written consent of Boston Biochem, Inc. "Boston Biochem" is a Trademark of Boston Biochem, Inc., 840 Memorial Drive, Cambridge, MA 02139.