

An R&D Systems Company

MATERIAL DATA SHEET

Di-Ubiquitin/Ub2, 6-DCA-linked, human recombinant Cat. # UCD-10

Dichloroacetone (DCA) linked forms of Ub₂ are chemically linked via a DCA linker using cysteine introduced into the ubiquitin sequence at the appropriate residues to mimic a native isopeptide linkage. Reports have demonstrated that the DCA linkage is stable and cannot be processed by deubiquitinating enzymes. Additionally, the orientation of the two ubiquitins in DCA-linked Ub₂ is preserved in relation to di-ubiquitin containing a native isopeptide linkage. This makes DCA-linked Ub₂ useful for characterizing polyubiquitin-binding proteins in the absence of added deubiquitinase inhibitors.

Product Information

Quantity: 100 μg, lyophilized powder

Solubility: Aqueous solutions $\geq 1 \text{ mg/ml}$

MW: 17 kDa

Purity: > 95% by SDS-PAGE and HPLC

Use & Storage

Use: Typical concentrations will depend on specific assay conditions and method of

detection.

Storage: Store at -80°C. Avoid multiple freeze/thaw cycles.

Literature

References: Yin L., et al. (2000) <u>Biochemistry</u> **39**: 10001-10010

Messick T., *et al.* (2008) <u>J Biol Chem.</u> **283**: 11038-11049 Carlile, C., *et al.* (2009) <u>J Biol Chem.</u> **284**: 29326-29334

For Laboratory Research Use Only, Not For Use in Humans

Rev: 12/06/2012

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-241-7072 FAX: 617-492-3565 www.bostonbiochem.com