

Lot # XXXXX

BostonBiochem®

An R&D Systems Company

MATERIAL DATA SHEET

Di-Ubiquitin/Ub₂, 29-DCA-linked, human recombinant

Cat. # UCD-80

Dichloroacetone (DCA) forms of di-ubiquitin (Ub₂) are chemically linked via a DCA linker using cysteine introduced into the ubiquitin sequence at the appropriate residues to mimic a native isopeptide linkage. Reports have demonstrated that the DCA linkage is stable and cannot be processed by deubiquitinases. Additionally, the orientation of the two ubiquitins in DCA-linked Ub₂ is preserved in relation to di-ubiquitin containing a native isopeptide linkage. This makes DCA-linked Ub₂ useful for characterizing polyubiquitin-binding proteins in the absence of added deubiquitinase inhibitors.

Product Information

Quantity:	100 µg, lyophilized powder
Solubility:	Aqueous solutions ≥ 1 mg/ml
MW:	17 kDa
Purity:	> 95% by SDS-PAGE and HPLC

Use & Storage

Use:	Typical concentrations will depend on specific assay conditions and method of detection.
Storage:	Resuspended material at -20°C. Avoid multiple freeze/thaw cycles.

Literature

References:	Yin L., <i>et al.</i> (2000) <u>Biochemistry</u> 39 : 10001-10010 Messick T., <i>et al.</i> (2008) <u>J Biol Chem.</u> 283 : 11038-11049 Carlile, C., <i>et al.</i> (2009) <u>J Biol Chem.</u> 284 : 29326-29334
--------------------	---

For Laboratory Research Use Only, Not For Use in Humans

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565
www.bostonbiochem.com

The contents of this datasheet (unless otherwise noted) are Copyright © 2008 Boston Biochem, Inc. All rights reserved. Duplication in whole or in part is strictly prohibited without the express written consent of Boston Biochem, Inc. "Boston Biochem" is a Trademark of Boston Biochem, Inc., 840 Memorial Drive, Cambridge, MA 02139.