

---

## MATERIAL DATA SHEET

---

### Recombinant Human Pro NEDD8

#### Cat. # UL-810

Neural Precursor Cell Expressed Developmentally Downregulated Gene 8 (NEDD8), also known as Related to Ubiquitin 1 (Rub1), is a 6-8 kDa member of the Ubiquitin family of proteins. Human pro-NEDD8 is 81 amino acids (aa) in length. Pro-NEDD8 contains one Ubiquitin-like domain (aa 1-65) and a 4-5 aa C-terminal propeptide (1,2). Cleavage of the propeptide exposes a C-terminal glycine residue that is used to generate a glycine-lysine intermolecular bond. Mature human NEDD8 shows 100% aa identity to NEDD8 from mouse, rat, and canine. Human NEDD8 is activated by a distinct NEDD8-activating (E1) enzyme, a heterodimeric complex composed of APPBP1 and UBA3 subunits (3). Activated NEDD8 is subsequently transferred to the UBE2M/Ubc12 or UBE2F NEDD8-conjugating (E2) enzymes (4). Through a process termed neddylation, the ROC1/Rbx1 RING Finger E3 ligase transfers NEDD8 to specific substrates (5). For example, transfer of NEDD8 to the Cullin scaffold activates members of the Cullin-RING Ligase (CRL) family of Ubiquitin ligases (E3) (6-8). CRL-induced ubiquitination is known to regulate proteins that are important for cell cycle progression and cell survival (9,10). Physiologically, NEDD8 plays a critical regulatory role in cell proliferation and dysregulation of the NEDD8 pathway has been associated with several cancer pathologies (11,12).

#### Product Information

<b>Quantity:</b>	1 mg
<b>Source:</b>	<i>E. coli</i> -derived human NEDD8 protein Accession # Q15843
<b>Stock:</b>	X mg/ml (X $\mu$ M) in 20 mM HEPES pH 8.0, 50 mM NaCl, 1 mM DTT
<b>Purity:</b>	>95%, by SDS-PAGE under reducing conditions and visualized by Colloidal Coomassie® Blue stain.

## Use & Storage

**Use:** Recombinant Human Pro-NEDD8 CANNOT be conjugated to substrate proteins via the subsequent actions of a NEDD8-activating (E1) enzyme, a NEDD8-conjugating (E2) enzyme, and a NEDD8 ligase (E3). Reaction conditions will need to be optimized for each specific application.

**Storage:** Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 48 months from date of receipt, -70 °C as supplied.
- 3 months, -70 °C under sterile conditions after opening.

## Literature

### References:

1. Kumar, S. *et al.* (1993) *Biochem. Biophys. Res. Commun.* **195**:393.
2. Kamitani, T. *et al.* (1997) *J. Biol. Chem.* **272**:28557.
3. Huang, D.T. & B.A. Schulman (2005) *Methods Enzymol.* **398**:9.
4. Gong, L. & E.T. Yeh (1999) *J. Biol. Chem.* **274**:12036.
5. Ohki, Y. *et al.* (2009) *Biochem. Biophys. Res. Commun.* **381**:443.
6. Petroski, M.D. & R.T. Deshares (2005) *Nat. Rev. Mol. Cell Biol.* **6**:9.
7. Hjerpe, R. *et al.* (2012) *Biochem. J.* **441**:927.
8. Calabrese, M.F. *et al.* (2011) *Nat. Struct. Mol. Biol.* **18**:947.
9. Tateishi, K. *et al.* (2001) *J. Cell. Biol.* **155**:571.
10. Ohh, M. *et al.* (2002) *EMBO Rep.* **3**:177.
11. Luo, Z. *et al.* (2012) *Cancer Res.* **72**:3360.
12. Soucy, T.A. *et al.* (2010) *Genes Cancer* **1**:708.

***For research use only. Not for use in humans.***