
MATERIAL DATA SHEET

Recombinant Human His6 NEDD8**Cat. # UL-813**

Neural Precursor Cell Expressed Developmentally Downregulated Gene 8 (NEDD8), also known as Related to Ubiquitin 1 (Rub1), is a 6-8 kDa member of the Ubiquitin family of proteins. Human pro-NEDD8 is 81 amino acids (aa) in length. Pro-NEDD8 contains one Ubiquitin-like domain (aa 1-65) and a 4-5 aa C-terminal propeptide (1,2). Cleavage of the propeptide exposes a C-terminal glycine residue that is used to generate a glycine-lysine intermolecular bond. Mature human NEDD8 shows 100% aa identity to NEDD8 from mouse, rat, and canine. Human NEDD8 is activated by a distinct NEDD8-activating (E1) enzyme, a heterodimeric complex composed of APPBP1 and UBA3 subunits (3). Activated NEDD8 is subsequently transferred to the UBE2M/Ubc12 or UBE2F NEDD8-conjugating (E2) enzymes (4). Through a process termed neddylation, the ROC1/Rbx1 RING Finger E3 ligase transfers NEDD8 to specific substrates (5). For example, transfer of NEDD8 to the Cullin scaffold activates members of the Cullin-RING Ligase (CRL) family of Ubiquitin ligases (E3) (6-8). CRL-induced ubiquitination is known to regulate proteins that are important for cell cycle progression and cell survival (9,10). Physiologically, NEDD8 plays a critical regulatory role in cell proliferation and dysregulation of the NEDD8 pathway has been associated with several cancer pathologies (11,12).

Product Information

Quantity:	500 µg
MW:	9.5 kDa
Source:	<i>E. coli</i> -derived Contains a 6-His tag Accession # Q15843
Stock:	Supplied as a solution in Tris, NaCl and Glycerol.
Purity:	>95%, by SDS-PAGE under reducing conditions and visualized by Colloidal Coomassie® Blue stain.

Use & Storage

Use: Recombinant Human His6-NEDD8 can be conjugated to substrate proteins via the subsequent actions of a NEDD8-activating (E1) enzyme, a NEDD8-conjugating (E2) enzyme, and a NEDD8 ligase (E3). Reaction conditions will need to be optimized for each specific application. We recommend an initial Recombinant Human His6-NEDD8 concentration of 5-20 μ M.

Storage: Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -70 °C as supplied.
- 3 months, -70 °C under sterile conditions after opening.

Literature

References:

1. Kumar, S. *et al.* (1993) *Biochem. Biophys. Res. Commun.* **195**:393.
2. Kamitani, T. *et al.* (1997) *J. Biol. Chem.* **272**:28557.
3. Huang, D.T. & B.A. Schulman (2005) *Methods Enzymol.* **398**:9.
4. Gong, L. & E.T. Yeh (1999) *J. Biol. Chem.* **274**:12036.
5. Ohki, Y. *et al.* (2009) *Biochem. Biophys. Res. Commun.* **381**:443.
6. Petroski, M.D. & R.T. Deshares (2005) *Nat. Rev. Mol. Cell Biol.* **6**:9.
7. Hjerpe, R. *et al.* (2012) *Biochem. J.* **441**:927.
8. Calabrese, M.F. *et al.* (2011) *Nat. Struct. Mol. Biol.* **18**:947.
9. Tateishi, K. *et al.* (2001) *J. Cell. Biol.* **155**:571.
10. Ohh, M. *et al.* (2002) *EMBO Rep.* **3**:177.
11. Luo, Z. *et al.* (2012) *Cancer Res.* **72**:3360.
12. Soucy, T.A. *et al.* (2010) *Genes Cancer* **1**:708.

For research use only. Not for use in humans.