

Product Name: (S)-(-)-5-Iodowillardiine


Catalog No.: 0307

Batch No.: 4

CAS Number: 140187-25-3

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₇H₈IN₃O₄
Batch Molecular Weight: 325.06
Physical Appearance: White solid
Solubility: 1.1eq. NaOH to 25 mM
Storage: Store at RT
Batch Molecular Structure:



2. ANALYTICAL DATA

TLC: R_f = 0.29 (Pyridine:Acetic acid:Water:Butanol [3:8:11:33])
Melting Point: At 238°C
HPLC: Shows 98% purity
Chiral HPLC: Shows >98% purity
¹H NMR: Consistent with structure
Optical Rotation: [α]_D = +15 (Concentration = 0.5, Solvent = 6N HCl)

Microanalysis:

	Carbon	Hydrogen	Nitrogen
Theoretical	25.86	2.48	12.93
Found	25.81	2.63	12.8

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: (S)-(-)-5-Iodowillardiine

Catalog No.: 0307

Batch No.: 4

CAS Number: 140187-25-3

Description:

Demonstrates high affinity for the kainate receptor subtype hGluK1 (formerly hGluR5) ($K_i = 0.24$ nM) and 600-4000-fold selectivity over both the AMPA receptor subtypes and the homomeric kainate receptor hGluK2 (formerly hGluR6). Please refer to IUPHAR Guide to Pharmacology for the most recent naming conventions.

Physical and Chemical Properties:


Batch Molecular Formula: C₇H₈IN₃O₄

Batch Molecular Weight: 325.06

Physical Appearance: White solid

Minimum Purity: >98%

Batch Molecular Structure:



Storage: Store at RT

CAUTION - This product is light sensitive and we recommend that the solid material and any solutions obtained are protected from exposure to light.

Solubility & Usage Info:

1.1eq. NaOH to 25 mM

Solutions of this product of up to 50mM (14mg/ml) are best prepared using 1eq. of NaOH solution with gentle warming if necessary (avoid excessive heating). Occasionally a slight excess - up to 15% - of NaOH may be advantageous. Aqueous solutions of pH7-9 are stable for several days; they should be kept frozen (-20°C) when not in use

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Swanson et al (1998) Kainate receptors exhibit differential sensitivities to (S)-5-iodowillardiine. *Mol.Pharmacol.* **53** 942. PMID: 9584222.

Jane et al (1997) Synthesis of willardiine and 6-azawillardiine analogs: pharmacological characterization on cloned homomeric human AMPA and kainate receptor subtypes. *J.Med.Chem.* **40** 3645. PMID: 9357531.

Thompson et al (1996) Depolarising effects of certain derivatives of (S) willardiine upon *in vitro* neonatal rat dorsal roots. *Br.J.Pharmacol.* **117** 331P.

Wong et al (1994) Willardiines differentiate agonist binding sites for kainate-versus AMPA-preferring glutamate receptors in DRG and hippocampal neurones. *J.Neurosci.* **14** 3881. PMID: 7515954.

Patneau et al (1992) Activation and desensitization of AMPA/kainate receptors by novel derivatives of willardiine. *J.Neurosci.* **12** 595. PMID: 1371315.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel:+1 612 379 2956