

# **Certificate of Analysis**

Print Date: Oct 21st 2024

www.tocris.com

Product Name: (+)-MK 801 maleate Catalog No.: 0924 Batch No.: 14

CAS Number: 77086-22-7 EC Number: 278-614-5

IUPAC Name: (5S,10R)-(+)-5-Methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5,10-imine maleate

#### 1. PHYSICAL AND CHEMICAL PROPERTIES

**Batch Molecular Formula:**  $C_{16}H_{15}N.C_4H_4O_4$ .

**Batch Molecular Weight:** 337.37 **Physical Appearance:** White solid

**Solubility:** water to 10 mM with gentle warming

DMSO to 100 mM

Storage: Store at RT

**Batch Molecular Structure:** 

HN Me .C<sub>d</sub>H<sub>d</sub>O<sub>d</sub>

2. ANALYTICAL DATA

HPLC: Shows 99.4% purity
Chiral HPLC: Shows 99.8% purity

<sup>1</sup>H NMR: Consistent with structure Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 71.2 5.68 4.15 Found 71.22 5.6 4.15

www.tocris.com/distributors Tel:+1 612 379 2956



# **Product Information**

Print Date: Oct 21st 2024

www.tocris.com

Catalog No.: 0924 14 **Product Name:** (+)-MK 801 maleate

CAS Number: 77086-22-7 EC Number: 278-614-5

**IUPAC Name:** (5S,10R)-(+)-5-Methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5,10-imine maleate

#### **Description:**

(+)-MK 801 maleate is a high affinity (K<sub>i</sub> = 37.2 nM), selective and non-competitive NMDA receptor antagonist. (+)-MK 801 maleate acts by binding to a site located within the NMDA associated ion channel and thus prevents Ca2+ flux. It is an effective anti-ischemic agent in several animal models. (+)-MK 801 maleate increases motor activity in rat models. (+)-MK 801 maleate inhibits proliferation and increases apoptosis in hippocampal neural stem cells (NSCs). (+)-MK 801 maleate induces schizophrenia-like symptoms in rodents. (-)-enantiomer also available.


#### **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>16</sub>H<sub>15</sub>N.C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>.

Batch Molecular Weight: 337.37 Physical Appearance: White solid

Minimum Purity: ≥99%

#### **Batch Molecular Structure:**



Storage: Store at RT

### Solubility & Usage Info:

water to 10 mM with gentle warming DMSO to 100 mM

## Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. \*Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

#### References:

Ding et al (2018) Effect of NMDA on proliferation and apoptosis in hippocampal neural stem cells treated with MK-801. Exp.Ther.Med. 16 1137. PMID: 30116364.

Carey et al (1998) Effects of dizocilpine (MK-801) on motor activity and memory. Psychopharmacology 137 241. PMID: 9683001.

Zajaczkowski et al (1997) Uncompetitive NMDA receptor antagonists attenuate NMDA-induced impairment of passive avoidance learning and LTP. Neuropharmacology 36 961. PMID: 9257940.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use