

# **Certificate of Analysis**

Print Date: Jan 6th 2022

www.tocris.com

Product Name: Endomorphin-2 Catalog No.: 1056 Batch No.: 11

CAS Number: 141801-26-5

## 1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{32}H_{37}N_5O_5$ Batch Molecular Weight:571.65

Physical Appearance: White lyophilised solid

Net Peptide Content: 82%
Counter Ion: TFA

**Solubility:** Soluble to 0.40 mg/ml in water

Storage: Store at -20°C

Peptide Sequence: Tyr-Pro-Phe-Phe-NH<sub>2</sub>

2. ANALYTICAL DATA

**HPLC:** Shows 97.7% purity

Mass Spectrum: Consistent with structure

3. AMINO ACID ANALYSIS DATA

Lys		
Met		
Phe	2.00	2.00
Pro	1.00	1.01
Ser		
Thr		
Trp		
Tyr	1.00	0.99
Val		
	Met Phe Pro Ser Thr Trp Tyr	Met Phe 2.00 Pro 1.00 Ser Thr Trp Tyr 1.00

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use



# **Product Information**

Print Date: Jan 6th 2022

www.tocris.com

Product Name: Endomorphin-2 Catalog No.: 1056 Batch No.: 11

CAS Number: 141801-26-5

### **Description:**

Endomorphin-2 is a endogenous peptide with extremely high affinity and selectivity for  $\mu$ -opioid receptors (with  $K_i$  values of 0.69, 9233 and 5240 nM for  $\mu$ ,  $\delta$  and  $\kappa$  respectively).

### **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>32</sub>H<sub>37</sub>N<sub>5</sub>O<sub>5</sub> Batch Molecular Weight: 571.65

Physical Appearance: White lyophilised solid

#### **Peptide Sequence:**

Tyr-Pro-Phe-Phe-NH<sub>2</sub>

Storage: Store at -20°C

#### Solubility & Usage Info:

Soluble to 0.40 mg/ml in water

Net Peptide Content: 82% (Remaining weight made up of

counterions and residual water).

**Counter Ion: TFA** 

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such Cys, Met,Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2  $\mu$ m filter to remove potential bacterial contamination whenever possible.

#### References:

**Goldberg** *et al* (1998) Pharmacological characterization of endomorphin-1 and endomorphin-2 in mouse brain. J.Pharmacol.Exp.Ther. **286** 1007. PMID: 9694962.

Harrison et al (1998) Differential effects of endomorphin-1, endomorphin-2, and tyr-W-MIF-1 on activation of G-proteins in SH-SY5Y human neuroblastoma membranes. Peptides 19 749. PMID: 9622031.

**Chapman** *et al* (1997) Distinct inhibitory effects of spinal endomorphin-1 and endomorphin-2 on evoked dorsal horn neuronal responses in the rat. Br.J.Pharmacol. *122* 1537. PMID: 9422796.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Tel: +86 (21) 52380373