

Product Name: Endothelin 2 (human)

Catalog No.: 1164

Batch No.: 15

CAS Number: 123562-20-9

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₁₁₅H₁₆₀O₃₂N₂₆S₄
Batch Molecular Weight: 2547
Physical Appearance: White lyophilised solid
Counter Ion: TFA
Solubility: Soluble to 1 mg/ml in DMSO
Storage: Store at -20°C
Peptide Sequence:

```

  Cys-Ser-Cys-Ser-Ser-Trp-Leu-Asp-Lys-Glu-
  Cys-Val-Tyr-Phe-Cys-His-Leu-Asp-Ile-Ile-Trp
  
```

2. ANALYTICAL DATA

HPLC: Shows 96.9% purity
Mass Spectrum: Consistent with structure

3. AMINO ACID ANALYSIS DATA

Amino Acid		Theoretical	Actual	Amino Acid		Theoretical	Actual
Ala				Lys	1.00	1.01	
Arg				Met			
Asx	2.00	2.02	Phe	1.00	1.02		
Cys	4.00	1.95	Pro				
Glx	1.00	1.00	Ser	3.00	2.10		
Gly			Thr				
His	1.00	1.00	Trp	2.00	0.05		
Ile	2.00	1.73	Tyr	1.00	0.93		
Leu	2.00	1.98	Val	1.00	1.05		

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: Endothelin 2 (human)

Catalog No.: 1164

15

CAS Number: 123562-20-9

Description:

Endothelin 2 (human) is an endogenous peptide found predominately in the kidney and intestine. Displays similar selectivity for ET_A and ET_B endothelin receptors.


Physical and Chemical Properties:

Batch Molecular Formula: C₁₁₅H₁₆₀O₃₂N₂₆S₄

Batch Molecular Weight: 2547

Physical Appearance: White lyophilised solid

Peptide Sequence:



Storage: Store at -20°C

Solubility & Usage Info:

Soluble to 1 mg/ml in DMSO

This product is supplied in lyophilized form. It may appear as a solid, gel or film and be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Counter Ion: TFA

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such as Cys, Met, Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 µm filter to remove potential bacterial contamination whenever possible.

References:

Servitja *et al* (1998) Involvement of ET_A and ET_B receptors in the activation of phospholipase D by endothelins in cultured rat cortical astrocytes. *Br.J.Pharmacol.* **124** 1728. PMID: 9756390.

Inoue *et al* (1989) The human endothelin family: three structurally and pharmacologically distinct isopeptides predicted by three separate genes. *Proc.Natl.Acad.Sci.U.S.A.* **86** 2863. PMID: 2649896.

Yanagisawa and Masaski (1989) Molecular biology and biochemistry of the endothelins. *TiPS* **10** 374. PMID: 2690429.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel:+1 612 379 2956