



Certificate of Analysis

www.tocris.com

Product Name: Cytochalasin D Catalog No.: 1233 Batch No.: 12

CAS Number: 22144-77-0 EC Number: 244-804-1

IUPAC Name: (7S,13E,16S,18R,19E,21R)-21-(Acetyloxy)-7,18-dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13,19-

triene-1,17-dione

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{30}H_{37}NO_6$.Batch Molecular Weight:507.63Physical Appearance:White solid

Solubility: ethanol to 5 mg/ml with gentle warming

DMSO to 25 mg/ml

Storage: Store at -20°C

Batch Molecular Structure:

2. ANALYTICAL DATA

HPLC: Shows 98.6% purity

¹H NMR: Consistent with structure Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 70.98 7.35 2.76 Found 71.96 7.47 2.84

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use



Product Information

Print Date: Aug 11th 2023

www.tocris.com

Product Name: Cytochalasin D Catalog No.: 1233 12

CAS Number: 22144-77-0 EC Number: 244-804-1

IUPAC Name: (7S,13E,16S,18R,19E,21R)-21-(Acetyloxy)-7,18-dihydroxy-16,18-dimethyl-10-phenyl[11]cytochalasa-6(12),13,19-

triene-1,17-dione

Description:

Cytochalasin D is a potent disruptor of actin filament function. Alters tight junction permeability. Unlike cytochalasin B (Cat. No. 5474), does not inhibit monosaccharide transport across the plasma membrane.

Physical and Chemical Properties:

Batch Molecular Formula: $C_{30}H_{37}NO_6$. Batch Molecular Weight: 507.63 Physical Appearance: White solid

Minimum Purity: ≥95%

Batch Molecular Structure:

Storage: Store at -20°C

Solubility & Usage Info:

ethanol to 5 mg/ml with gentle warming DMSO to 25 mg/ml

When purched as a 1mg unit, this product is supplied as a lyophilized solid and may be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. *Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Abedi and Zachary (1998) Cytochalasin D stimulation of tyrosine phosphorylation and phosphotyrosine-associated kinase activity in vascular smooth muscle cells. Biochem.Biophys.Res.Commun. **245** 646. PMID: 9588169.

Stevenson and Begg (1994) Concentration-dependent effects of cytochalasin D on tight junctions and actin filaments in MDCK epithelial cells. J.Cell Sci. **107** 367. PMID: 8006058.

Carlier et al (1986) Interaction of cytochalasin D with actin filaments in the presence of ADP and ATP. J.Biol.Chem. 261 2041. PMID: 3944126.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use