

**Product Name:** Rapamycin

**Catalog No.:** 1292


**Batch No.:** 10

CAS Number: 53123-88-9

IUPAC Name: (3*S*,6*R*,7*E*,9*R*,10*R*,12*R*,14*S*,15*E*,17*E*,19*E*,21*S*,23*S*,26*R*,27*R*,34*aS*)-9,10,12,13,14,21,22,23,24,25,26,27,32,33,34,34*a*-Hexadecahydro-9,27-dihydroxy-3-[(1*R*)-2-[(1*S*,3*R*,4*R*)-4-hydroxy-3-methoxycyclohexyl]-1-methylethyl]-10,21-dimethoxy-6,8,12,14,20,26-hexamethyl-23,27-epoxy-3*H*-pyrido[2,1-*c*][1,4]oxaazacyclohentriacontine-1,5,11,28,29(4*H*,6*H*,31*H*)-pentone

**1. PHYSICAL AND CHEMICAL PROPERTIES**

**Batch Molecular Formula:** C<sub>51</sub>H<sub>79</sub>NO<sub>13</sub>  
**Batch Molecular Weight:** 914.18  
**Physical Appearance:** White solid  
**Solubility:** DMSO to 20 mM  
**Storage:** Store at -20°C  
**Batch Molecular Structure:**



**2. ANALYTICAL DATA**

**HPLC:** Shows 98.1% purity  
**Mass Spectrum:** Consistent with structure

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

<b>Product Name:</b>	<b>Rapamycin</b>	<b>Catalog No.: 1292</b>	<b>10</b>
CAS Number:	53123-88-9		
IUPAC Name:	(3 <i>S</i> ,6 <i>R</i> ,7 <i>E</i> ,9 <i>R</i> ,10 <i>R</i> ,12 <i>R</i> ,14 <i>S</i> ,15 <i>E</i> ,17 <i>E</i> ,19 <i>E</i> ,21 <i>S</i> ,23 <i>S</i> ,26 <i>R</i> ,27 <i>R</i> ,34 <i>aS</i> )-9,10,12,13,14,21,22,23,24,25,26,27,32,33,34,34a-Hexadecahydro-9,27-dihydroxy-3-[(1 <i>R</i> )-2-[(1 <i>S</i> ,3 <i>R</i> ,4 <i>R</i> )-4-hydroxy-3-methoxycyclohexyl]-1-methylethyl]-10,21-dimethoxy-6,8,12,14,20,26-hexamethyl-23,27-epoxy-3 <i>H</i> -pyrido[2,1- <i>c</i> ][1,4]oxaazacyclohentriacontine-1,5,11,28,29 (4 <i>H</i> ,6 <i>H</i> ,31 <i>H</i> )-pentone		

**Description:**


Rapamycin is an antifungal and immunosuppressant. Rapamycin is a specific inhibitor of mTOR (mammalian target of rapamycin); complexes with FKBP-12 and binds mTOR inhibiting its activity. Rapamycin inhibits interleukin-2-induced phosphorylation and activation of p70 S6 kinase and induces autophagy in yeast and mammalian cell lines. Rapamycin drives hPSC differentiation to mesendoderm and blood progenitor cells. Also used as a chemical dimerizer; rapamycin and GA<sub>3</sub>-AM chemically inducible dimerization systems are orthogonal. Inhibits MERS-CoV infection of Huh7 cells in a plaque reduction assay. Enhances lentiviral transduction of hematopoietic ... Please see product specific page on [www.tocris.com](http://www.tocris.com) for full description.

**Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>51</sub>H<sub>79</sub>NO<sub>13</sub>  
 Batch Molecular Weight: 914.18  
 Physical Appearance: White solid

**Minimum Purity:** ≥98%

**Batch Molecular Structure:**



**Storage:** Store at -20°C

**Solubility & Usage Info:**

DMSO to 20 mM

This product is supplied as a lyophilized solid and may be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

**Stability and Solubility Advice:**

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. \*Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

**SOLIDS:** Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

**SOLUTIONS:** We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

**References:**

**Galluzzi et al** (2017) Pharmacological modulation of autophagy: therapeutic potential and persisting obstacles. *Nat.Rev.Drug.Discov.* . PMID: 28529316 . Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**Nazareth et al** (2016) A multi-lineage screen reveals mTORC1 inhibition enhances human pluripotent stem cell mesendoderm and blood progenitor production. *Stem Cells* 34(12):3174-3187. PMID: 27132889.

**Kindler et al** (2015) Antiviral activity of ERK1/2 inhibitor SBKAT/mTOR inhibitor rapamycin for Middle East Respiratory Syndrome Coronavirus infection as identified by temporal genome analysis. *Antimicrob.Agents Chemother.* 59(10):3147-3154. PMID: 25967801.

biotechne.com    North America    China    Europe Middle East Africa    Rest of World  
 Tel: +1 610 279 2956    Tel: +86 (21) 52380373    Tel: +44 (1473) 209000    Tel: +91 80 4709 9200