

**Product Name:** Gap 27  
CAS Number: 198284-64-9

**Catalog No.:** 1476 **Batch No.:** 14

**1. PHYSICAL AND CHEMICAL PROPERTIES**

**Batch Molecular Formula:** C<sub>60</sub>H<sub>101</sub>N<sub>15</sub>O<sub>17</sub>  
**Batch Molecular Weight:** 1304.55  
**Physical Appearance:** White lyophilised solid  
**Counter Ion:** TFA  
**Solubility:** Soluble to 1 mg/ml in water  
**Storage:** Store at -20°C  
**Peptide Sequence:** Ser-Arg-Pro-Thr-Glu-Lys-Thr-Ile-Phe-Ile-Ile

**2. ANALYTICAL DATA**

**HPLC:** Shows 99.0% purity  
**Mass Spectrum:** Consistent with structure

**3. AMINO ACID ANALYSIS DATA**

Amino Acid		Theoretical	Actual	Amino Acid		Theoretical	Actual
Ala				Lys	1.00	0.99	
Arg	1.00	0.99	Met				
Asx			Phe	1.00	1.00		
Cys			Pro	1.00	0.99		
Glx	1.00	1.02	Ser	1.00	1.01		
Gly			Thr	2.00	2.06		
His			Trp				
Ile	3.00	2.14	Tyr				
Leu			Val				

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**Product Name:** Gap 27**Catalog No.:** 1476**14**

CAS Number: 198284-64-9

**Description:**

Gap 27 is a peptide derived from connexin 43 that is a selective gap junction blocker. Attenuates ACh-induced arterial relaxation and reduces K<sup>+</sup>-mediated smooth muscle repolarization in endothelium-intact vessels in vitro.

**Physical and Chemical Properties:**Batch Molecular Formula: C<sub>60</sub>H<sub>101</sub>N<sub>15</sub>O<sub>17</sub>

Batch Molecular Weight: 1304.55

Physical Appearance: White lyophilised solid

**Peptide Sequence:**

Ser-Arg-Pro-Thr-Glu-Lys-Thr-Ile-Phe-Ile-Ile

**Storage:** Store at -20°C**Solubility & Usage Info:**

Soluble to 1 mg/ml in water

This product is supplied in lyophilized form. It may appear as a solid, gel or film and be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

**Counter Ion:** TFA**Stability and Solubility Advice:**

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such as Cys, Met, Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 µm filter to remove potential bacterial contamination whenever possible.

**References:**

**Richards *et al*** (2001) Suppression of K<sup>+</sup>-induced hyperpolarization by phenylephrine in rat mesenteric artery: relevance to studies of endothelium-derived hyperpolarizing factor. *Br.J.Pharmacol.* **134** 1. PMID: 11522590.

**Ko *et al*** (2000) Biochemical and functional characterization of intercellular adhesion and gap junctions in fibroblasts. *Am.J.Physiol.Cell Physiol.* **279** C147. PMID: 10898726.

**Chaytor *et al*** (1998) Central role of heterocellular gap junctional communication in endothelium-dependent relaxations of rabbit arteries. *J.Physiol.* **508** 561. PMID: 9508817.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**bio-techne.com**

info@bio-techne.com

techsupport@bio-techne.com

**North America**

Tel: (800) 343 7475

**China**

info.cn@bio-techne.com

Tel: +86 (21) 52380373

**Europe Middle East Africa**

Tel: +44 (0)1235 529449

**Rest of World**

www.tocris.com/distributors

Tel:+1 612 379 2956