

Product Name: Demethylasterriquinone B1

Catalog No.: 1819


Batch No.: 3

CAS Number: 78860-34-1

IUPAC Name: 2-[2-(1,1-Dimethyl-2-propenyl)-1*H*-indol-3-yl]-3,6-dihydroxy-5-[7-(3-methyl-2-butenyl)-1*H*-indol-3-yl]-2,5-cyclohexadiene-1,4-dione

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula:	C ₃₂ H ₃₀ N ₂ O ₄
Batch Molecular Weight:	506.59
Physical Appearance:	Black solid
Solubility:	DMSO to 100 mM
Storage:	Desiccate at -20°C
Batch Molecular Structure:	



2. ANALYTICAL DATA

HPLC:	Shows 98.1% purity
¹H NMR:	Consistent with structure
Mass Spectrum:	Consistent with structure

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: Demethylasterriquinone B1

Catalog No.: 1819

Batch No.: 3

CAS Number: 78860-34-1

IUPAC Name: 2-[2-(1,1-Dimethyl-2-propenyl)-1H-indol-3-yl]-3,6-dihydroxy-5-[7-(3-methyl-2-butenyl)-1H-indol-3-yl]-2,5-cyclohexadiene-1,4-dione

Description:

Selective insulin receptor (IR) activator (EC₅₀ values are 3 - 6 μM for IRTK and 100 μM for IGF1R and EGFR). Increases IR β subunit tyrosine phosphorylation and activation of PI 3-kinase and Akt, but not ERK. Induces glucose uptake in adipocytes and skeletal muscle in vitro, without enhancing vascular proliferation. Binds GAPDH. Also activates Trk by interacting at a site distinct from the neurotrophin-binding site.

Physical and Chemical Properties:


Batch Molecular Formula: C₃₂H₃₀N₂O₄

Batch Molecular Weight: 506.59

Physical Appearance: Black solid

Minimum Purity: >98%

Batch Molecular Structure:



Storage: Desiccate at -20°C

Solubility & Usage Info:

DMSO to 100 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Weber et al (2000) A novel insulin mimetic without a proliferative effect on vascular smooth muscle cells. *J.Vasc.Surg.* **32** 1118. PMID: 11107083.

Salituro et al (2001) Discovery of a small molecule insulin receptor activator. *Recent Prog.Horm.Res.* **56** 107. PMID: 11237209.

Wilkie et al (2001) The non-peptidyl fungal metabolite L-783,281 activates TRK neurotrophin receptors. *J.Neurochem.* **78** 1135. PMID: 11553687.

Webster et al (2003) Signaling effects of demethylasterriquinone B1, a selective insulin receptor modulator. *Chembiochem* **4** 379. PMID: 12740809.

Kim et al (2007) Glyceraldehyde 3-phosphate dehydrogenase is a cellular target of the insulin mimic demethylasterriquinone B1. *J.Med.Chem.* **50** 3423. PMID: 17595071.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel: +1 612 379 2956