



Certificate of Analysis

www.tocris.com

Product Name: Simvastatin Catalog No.: 1965 Batch No.: 2

CAS Number: 79902-63-9

IUPAC Name: (1S,3R,7S,8S,8aR)-1,2,3,7,8,8a-Hexahydro-3,7-dimethyl-8-[2-[(2R,4R)-tetrahydro-4-hydroxy-6-oxo-2*H*-pyran-2-yl]

ethyl]-1-naphthalenyly-2,2-dimethyl butanoate

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{25}H_{38}O_5$ Batch Molecular Weight: 418.57

Physical Appearance: White crystalline solid

Solubility: DMSO to 50 mM

ethanol to 75 mM

Storage: Desiccate at -20°C

Batch Molecular Structure:

2. ANALYTICAL DATA

HPLC: Shows >98.2% purity

¹H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 71.74 9.15 Found 71.64 9.25

Tel: +44 (0)1235 529449 www.tocris.com/distributors Tel:+1 612 379 2956



Product Information

Print Date: Dec 14th 2021

www.tocris.com

Product Name: Simvastatin Catalog No.: 1965 Batch No.: 2

CAS Number: 79902-63-9

IUPAC Name: (1S,3R,7S,8S,8aR)-1,2,3,7,8,8a-Hexahydro-3,7-dimethyl-8-[2-[(2R,4R)-tetrahydro-4-hydroxy-6-oxo-2*H*-pyran-2-yl]

ethyl]-1-naphthalenyly-2,2-dimethyl butanoate

Description:

Simvastatin is a HMG-CoA reductase inhibitor; decreases levels of low density lipoprotein. Has multiple biological effects including bone formation stimulation, inhibition of smooth muscle cell proliferation and migration, induction of ferroptosis, and anticancer and anti-inflammatory activity. Inactive lactone prodrug of simvastatin hydroxy acid, naturally bioactivated in vivo following oral administration. Simvastatin acts synergistically with ABL allosteric inhibitors GNF 5 (Cat. No. 1965) to enhance apoptosis of metastatic lung cancer cells in mice.

Physical and Chemical Properties:

Batch Molecular Formula: C₂₅H₃₈O₅ Batch Molecular Weight: 418.57

Physical Appearance: White crystalline solid

Minimum Purity: ≥98%

Batch Molecular Structure:

Storage: Desiccate at -20°C

Solubility & Usage Info:

DMSO to 50 mM ethanol to 75 mM

PLEASE NOTE - Simvastatin (SV) is an inactive lactone prodrug of simvastatin hydroxy acid (SVA) the active form of the compound. Simvastatin administered orally is naturally bioactivated by the liver and requires no further modification. If simvastatin is to be administered by another route or used *in vitro* then it requires manual activation by treatment with NaOH. To activate the compound dissolve 50mg in 1ml of warm (50°C) ethanol and add 0.813ml of 1N NaOH. Leave for 30 minutes to allow conversion of the simvastatin to the active acid form, the compound may be stored at -20°C in this format for up to 1 month. Adjust pH to 7.2 with small quantities of 1N HCl prior to use.

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Luttman et al (2021) ABL allosteric inhibitors synergize with statins to enhance apoptosis of metastatic lung cancer cells. Cell Rep. 37 109880. PMID: 34706244.

Kaushal et al (2003) Potential anticancer effects of STAT: fact or fiction? Endothelium 10 49. PMID: 12699077.

Reinoso et al (2002) Preclinical pharmacokinetics of STAT. Methods Find.Exp.Clin.Pharmacol. 24 593. PMID: 12616706.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use