

Product Name: L-161,982

Catalog No.: 2514


Batch No.: 2

CAS Number: 147776-06-5

IUPAC Name: *N*-[[4'-[[3-Butyl-1,5-dihydro-5-oxo-1-[2-(trifluoromethyl)phenyl]-4*H*-1,2,4-triazol-4-yl]methyl][1,1'-biphenyl]-2-yl]sulfonyl]-3-methyl-2-thiophenecarboxamide

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula:	C ₃₂ H ₂₉ F ₃ N ₄ O ₄ S ₂
Batch Molecular Weight:	654.72
Physical Appearance:	White solid
Solubility:	DMSO to 100 mM
Storage:	Desiccate at +4°C
Batch Molecular Structure:	



2. ANALYTICAL DATA

TLC:	R _f = 0.3 (Dichloromethane:Methanol [9:1])
HPLC:	Shows 99.4% purity
¹H NMR:	Consistent with structure
Mass Spectrum:	Consistent with structure
Microanalysis:	

	Carbon	Hydrogen	Nitrogen
Theoretical	58.7	4.46	8.55
Found	58.46	4.39	8.52

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: L-161,982

Catalog No.: 2514

Batch No.: 2

CAS Number: 147776-06-5

IUPAC Name: *N*-[[4'-[[3-Butyl-1,5-dihydro-5-oxo-1-[2-(trifluoromethyl)phenyl]-4*H*-1,2,4-triazol-4-yl]methyl][1,1'-biphenyl]-2-yl]sulfonyl]-3-methyl-2-thiophenecarboxamide

Description:

EP₄ receptor antagonist that is selective over all other members of the prostanoid receptor family (K_i values are 0.024, 0.71, 1.90, 5.10, 5.63, 6.74, 19 and 23 μM for human EP₄, TP, EP₃, DP, FP, IP, EP₁ and EP₂ receptors respectively). Suppresses PGE₂-induced bone formation in rats and prevents the nociceptive response induced by misoprostol in formalin-injected mice.

Physical and Chemical Properties:


Batch Molecular Formula: C₃₂H₂₉F₃N₄O₄S₂

Batch Molecular Weight: 654.72

Physical Appearance: White solid

Minimum Purity: >99%

Batch Molecular Structure:



Storage: Desiccate at +4°C

Solubility & Usage Info:

DMSO to 100 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Balzary et al (2006) Lipopolysaccharide induces epithelium- and prostaglandin E₂-dependent relaxation of mouse isolated trachea through activation of cyclooxygenase (COX)-1 and COX-2. *J.Pharmacol.Exp.Ther.* **317** 806. PMID: 16464966.

Oliva et al (2006) Role of periaqueductal grey prostaglandin receptors in formalin-induced hyperalgesia. *Eur.J.Pharmacol.* **530** 40. PMID: 16360148.

Machwate et al (2001) Prostaglandin receptor EP₄ mediates the bone anabolic effects of PGE₂. *Mol.Pharmacol.* **60** 36. PMID: 11408598.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel:+1 612 379 2956