

Product Name: Tunicamycin

Catalog No.: 3516

Batch No.: 8

CAS Number: 11089-65-9

IUPAC Name: Tunicamycin from *Streptomyces* sp.

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₃₉H₆₄N₄O₁₆ (tunicamycin C, n=10)


Batch Molecular Weight: 844.95 (tunicamycin C, n=10)

Physical Appearance: Off-white solid

Solubility: DMSO to 50 mM

Storage: Store at +4°C

Batch Molecular Structure:



2. ANALYTICAL DATA

Composition by HPLC:

Tunicamycin A	4.92%
Tunicamycin B	32.26%
Tunicamycin C	42.12%
Tunicamycin D	20.32%

TLC: R_f = 0.31 (Butanol:Acetic acid:Water [60:15:25])

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: Tunicamycin

Catalog No.: 3516

Batch No.: 8

CAS Number: 11089-65-9

IUPAC Name: Tunicamycin from *Streptomyces* sp.

Description:

Antibiotic; inhibits GlcNAc phosphotransferase (GPT). Blocks the formation of N-glycosidic linkages by inhibiting the first step in glycoprotein synthesis. Activity induces ER stress and causes G₁ arrest; can be used to induce autophagy. Tunicamycin contains four main components as follows: Homolog A, n=8, C₃₇H₆₀N₄O₁₆, molecular weight = 816.90 Homolog B, n=9, C₃₈H₆₂N₄O₁₆, molecular weight = 830.93 Homolog C, n=10, C₃₉H₆₄N₄O₁₆, molecular weight = 844.95 Homolog D, n=11, C₄₀H₆₆N₄O₁₆, molecular weight = 858.99 The composition of this product will vary from batch to batch and can be found on the relevant certificate of analysis. Please see product datasheet on www.tocris.com for full description.


Physical and Chemical Properties:

Batch Molecular Formula: C₃₉H₆₄N₄O₁₆ (tunicamycin C, n=10)

Batch Molecular Weight: 844.95

Physical Appearance: Beige solid

Batch Molecular Structure:



References:

Lauer *et al* (2009) Primary murine airway smooth muscle cells exposed to poly(I:C) or tunicamycin synthesize a leukocyte-adhesive hyaluronan matrix. *J.Biol.Chem.* **284** 5299. PMID: 19088077.

Duriez *et al* (2008) The hepatitis B virus precore protein is retrotransported from endoplasmic reticulum (ER) to cytosol through the ER-associated pathway. *J.Biol.Chem.* **283** 32352. PMID: 18805786.

Ding *et al* (2007) Differential effects of endoplasmic reticulum stress-induced autophagy on cell survival. *J.Biol.Chem.* **282** 4702. PMID: 17135238.

Storage: Store at +4°C

Solubility & Usage Info:

DMSO to 50 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel:+1 612 379 2956