

Certificate of Analysis

www.tocris.com

Product Name: ONO AE3 208

Catalog No.: 3565


Batch No.: 4

CAS Number: 402473-54-5

IUPAC Name: 4-Cyano-2-[[2-(4-fluoro-1-naphthalenyl)-1-oxopropyl]amino]benzenebutanoic acid

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula:	C ₂₄ H ₂₁ FN ₂ O ₃
Batch Molecular Weight:	404.43
Physical Appearance:	White solid
Solubility:	DMSO to 100 mM
Storage:	Store at -20°C
Batch Molecular Structure:	



2. ANALYTICAL DATA

HPLC:	Shows 99.9% purity
¹H NMR:	Consistent with structure
Mass Spectrum:	Consistent with structure

Microanalysis:	Carbon Hydrogen Nitrogen		
Theoretical	71.27	5.23	6.93
Found	70.62	5.24	6.74

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com
info@bio-techne.com
techsupport@bio-techne.com

North America
Tel: (800) 343 7475

China
info.cn@bio-techne.com
Tel: +86 (21) 52380373

Europe Middle East Africa
Tel: +44 (0)1235 529449

Rest of World
www.tocris.com/distributors
Tel: +1 612 379 2956

Product Name: ONO AE3 208

Catalog No.: 3565

4

CAS Number: 402473-54-5

IUPAC Name: 4-Cyano-2-[[2-(4-fluoro-1-naphthalenyl)-1-oxopropyl]amino]benzenebutanoic acid

Description:

ONO AE3 208 is a high affinity and selective EP₄ receptor antagonist (K_i values are 1.3, 30, 790 and 2400 nM for EP₄, EP₃, FP and TP receptors respectively). Displays no affinity for EP₁, EP₂, DP or IP receptors (K_i >10 μM). Inhibits PGE₂-induced IL-8 production in colonic epithelial caco-2 cells and attenuates PGE₂ inhibition of natural killer T cell activation. Suppresses recovery from experimentally-induced colitis and stimulates CD4⁺ T cell proliferation in C57BL/6 mice. Also reduces metastasis of mammary tumor cells in a murine model of breast cancer. Orally active. Please see product specific page on www.tocris.com for full description.

Physical and Chemical Properties:


Batch Molecular Formula: C₂₄H₂₁FN₂O₃

Batch Molecular Weight: 404.43

Physical Appearance: White solid

Minimum Purity: ≥98%

Batch Molecular Structure:



References:

Deng et al (2013) Exosome-like nanoparticles from intestinal mucosal cells carry prostaglandin E2 and suppress activation of liver NKT cells. *J.Immunol.* **190** 3579. PMID: 23467936.

Dey et al (2009) Prostaglandin E₂ couples through EP₄ prostanoid receptors to induce IL-8 production in human colonic epithelial cell lines. *Br.J.Pharmacol* **3** 475. PMID: 19175605.

Ma et al (2006) Prostaglandin E receptor EP₄ antagonism inhibits breast cancer metastasis. *Cancer Res.* **66** 2923. PMID: 16540639.

Storage: Store at -20°C

Solubility & Usage Info:

DMSO to 100 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com
techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com
Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors
Tel:+1 612 379 2956