


**Product Name:** RepSox  
**CAS Number:** 446859-33-2  
**IUPAC Name:** 2-(3-(6-Methylpyridine-2-yl)-1*H*-pyrazol-4-yl)-1,5-naphthyridine

**Catalog No.:** 3742      **Batch No.:** 10

**1. PHYSICAL AND CHEMICAL PROPERTIES**

**Batch Molecular Formula:** C<sub>17</sub>H<sub>13</sub>N<sub>5</sub>.  
**Batch Molecular Weight:** 287.32  
**Physical Appearance:** Pale yellow solid  
**Solubility:** DMSO to 100 mM  
 ethanol to 20 mM  
**Storage:** Store at +4°C  
**Batch Molecular Structure:**



**2. ANALYTICAL DATA**

**HPLC:** Shows 99.1% purity  
**<sup>1</sup>H NMR:** Consistent with structure  
**Mass Spectrum:** Consistent with structure

Microanalysis:	Carbon Hydrogen Nitrogen		
Theoretical	71.06	4.56	24.37
Found	70.65	4.56	24.37

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**Product Name:** RepSox

**Catalog No.:** 3742

**10**

CAS Number: 446859-33-2

IUPAC Name: 2-(3-(6-Methylpyridine-2-yl)-1H-pyrazol-4-yl)-1,5-naphthyridine

**Description:**

RepSox is a potent and selective inhibitor of the TGF- $\beta$  type I receptor/ALK5 (IC<sub>50</sub> values are 4 and 23 nM for TGF- $\beta$  type I receptor autophosphorylation and binding respectively). RepSox is selective for TGF- $\beta$  type I receptor over a range of kinases, including p38 MAPK, JNK1 and GSK3 (IC<sub>50</sub> > 16  $\mu$ M). Enhances the efficiency of cellular reprogramming; replaces Sox2 by inducing Nanog expression. RepSox synthesized to Ancillary Material Grade also available. For more information about how RepSox may be used, see our protocols: Highly Efficient Generation of CiPSCs from MEFs, Generation of  $\beta$  cells from hPSCs, Repro... Please see product specific page on www.tocris.com for full description.

**Physical and Chemical Properties:**


Batch Molecular Formula: C<sub>17</sub>H<sub>13</sub>N<sub>5</sub>.

Batch Molecular Weight: 287.32

Physical Appearance: Pale yellow solid

**Minimum Purity:**  $\geq$ 99%

**Batch Molecular Structure:**



**References:**

**Ichida *et al* (2009)** A small-molecule inhibitor of Tgf- $\beta$  signaling replaces Sox2 in reprogramming by inducing *Nanog*. *Cell Stem Cell* **5** 491. PMID: 19818703.

**Li *et al* (2009)** Generation of rat and human induced pluripotent stem cells by combining genetic reprogramming and chemical inhibitors. *Cell Stem Cell* **4** 16. PMID: 19097958.

**Gellibert *et al* (2004)** Identification of 1,5-naphthyridine derivatives as a novel series of potent and selective TGF- $\gamma$  type I receptor inhibitors. *J.Med.Chem.* **47** 4494. PMID: 15317461.

**Storage:** Store at +4°C

**Solubility & Usage Info:**

DMSO to 100 mM

ethanol to 20 mM

**Stability and Solubility Advice:**

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. \*Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

**SOLIDS:** Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

**SOLUTIONS:** We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**bio-techne.com**

info@bio-techne.com

techsupport@bio-techne.com

**North America**

Tel: (800) 343 7475

**China**

info.cn@bio-techne.com

Tel: +86 (21) 52380373

**Europe Middle East Africa**

Tel: +44 (0)1235 529449

**Rest of World**

www.tocris.com/distributors

Tel:+1 612 379 2956