

## Certificate of Analysis

[www.tocris.com](http://www.tocris.com)

**Product Name:** Pep2m, myristoylated

**Catalog No.:** 3801

**Batch No.:** 6

CAS Number: 1423381-07-0

### 1. PHYSICAL AND CHEMICAL PROPERTIES

**Batch Molecular Formula:** C<sub>63</sub>H<sub>118</sub>N<sub>18</sub>O<sub>14</sub>S  
**Batch Molecular Weight:** 1383.8  
**Physical Appearance:** White lyophilised solid  
**Net Peptide Content:** 74%  
**Counter Ion:** TFA  
**Solubility:** Soluble to 1 mg/ml in water  
**Storage:** Store at -20°C  
**Peptide Sequence:** Myr-Lys-Arg-Met-Lys-Val-Ala-Lys-Asn-Ala-Gln

### 2. ANALYTICAL DATA

**HPLC:** Shows 99% purity  
**Mass Spectrum:** Consistent with structure

### 3. AMINO ACID ANALYSIS DATA

Amino Acid		Theoretical	Actual	Amino Acid		Theoretical	Actual
Ala	2.00	1.90	Lys	3.00	3.09		
Arg	1.00	1.03	Met	1.00	0.98		
Asx	1.00	1.01	Phe				
Cys			Pro				
Glx	1.00	1.02	Ser				
Gly			Thr				
His			Trp				
Ile			Tyr				
Leu			Val	1.00	0.99		

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**bio-techne.com**

info@bio-techne.com

techsupport@bio-techne.com

**North America**

Tel: (800) 343 7475

**China**

info.cn@bio-techne.com

Tel: +86 (21) 52380373

**Europe Middle East Africa**

Tel: +44 (0)1235 529449

**Rest of World**

[www.tocris.com/distributors](http://www.tocris.com/distributors)

Tel: +1 612 379 2956

**Product Name:** Pep2m, myristoylated

**Catalog No.:** 3801

**Batch No.:** 6

CAS Number: 1423381-07-0

**Description:**

Cell-permeable, myristoylated form of pep2m (Cat. No. 1595). Peptide inhibitor of the interaction between the C-terminus of the GluA2 (AMPA receptor) subunit and N-ethylmaleimide-sensitive fusion protein (NSF), a protein that regulates AMPA receptor function. Reduces postsynaptic currents in CA1 neurons, AMPA-mediated currents in cultured hippocampal neurons and AMPA receptor surface expression.

**Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>63</sub>H<sub>118</sub>N<sub>18</sub>O<sub>14</sub>S

Batch Molecular Weight: 1383.8

Physical Appearance: White lyophilised solid

**Peptide Sequence:**

Myr-Lys-Arg-Met-Lys-Val-Ala-Lys-Asn-Ala-Gln

**Storage:** Store at -20°C

**Solubility & Usage Info:**

Soluble to 1 mg/ml in water

This product is supplied as a lyophilized solid and may be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

**Net Peptide Content:** 74% (Remaining weight made up of counterions and residual water).

**Counter Ion:** TFA

**Stability and Solubility Advice:**

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such as Cys, Met, Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 µm filter to remove potential bacterial contamination whenever possible.

**References:**

**Yao et al** (2008) PKMz maintains late long-term potentiation by N-ethylmaleimide-sensitive factor/GluR2-dependent trafficking of postsynaptic AMPA receptors. *J.Neurosci.* **28** 7820. PMID: 18667614.

**Luscher et al** (1999) Role of AMPA receptor cycling in synaptic transmission and plasticity. *Neuron* **24** 649. PMID: 10595516.

**Luthi et al** (1999) Hippocampal LTD expression involves a pool of AMPARs regulated by the NSF-GluR2 interaction. *Neuron* **24** 389. PMID: 10571232.

**Noel et al** (1999) Surface expression of AMPA receptors in hippocampal neurons is regulated by an NSF-dependent mechanism. *Neuron* **23** 365. PMID: 10399941.

**Nishimune et al** (1998) NSF binding to GluR2 regulates synaptic transmission. *Neuron* **21** 87. PMID: 9697854.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**bio-techne.com**

info@bio-techne.com

techsupport@bio-techne.com

**North America**

Tel: (800) 343 7475

**China**

info.cn@bio-techne.com

Tel: +86 (21) 52380373

**Europe Middle East Africa**

Tel: +44 (0)1235 529449

**Rest of World**

www.tocris.com/distributors

Tel:+1 612 379 2956