

**Product Name:** Deoxynivalenol

**Catalog No.:** 3976


**Batch No.:** 4

CAS Number: 51481-10-8

IUPAC Name: 3 $\alpha$ ,7 $\alpha$ ,15-Trihydroxy-12,13-epoxytrichothec-9-en-8-one

**1. PHYSICAL AND CHEMICAL PROPERTIES**

**Batch Molecular Formula:** C<sub>15</sub>H<sub>20</sub>O<sub>6</sub>  
**Batch Molecular Weight:** 296.32  
**Physical Appearance:** White solid  
**Solubility:** ethanol to 30 mM  
**Storage:** Store at +4°C  
**Batch Molecular Structure:**



**2. ANALYTICAL DATA**

**Melting Point:** At 155°C  
**HPLC:** Shows 100% purity

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**bio-techne.com**  
 info@bio-techne.com  
 techsupport@bio-techne.com

**North America**  
 Tel: (800) 343 7475

**China**  
 info.cn@bio-techne.com  
 Tel: +86 (21) 52380373

**Europe Middle East Africa**  
 Tel: +44 (0)1235 529449

**Rest of World**  
 www.tocris.com/distributors  
 Tel:+1 612 379 2956

**Product Name:** Deoxynivalenol

**Catalog No.:** 3976

**Batch No.:** 4

CAS Number: 51481-10-8

IUPAC Name: 3 $\alpha$ ,7 $\alpha$ ,15-Trihydroxy-12,13-epoxytrichothec-9-en-8-one

**Description:**

Tricothecene mycotoxin and potent protein synthesis inhibitor. Exhibits cytotoxic activity in vivo via the ribotoxic stress response. Induces p38-mediated gene expression and apoptosis in leukocytes; activity results in systemic expression of interleukin-6 (IL-6) and other proinflammatory cytokines. Also induces migration of NF- $\kappa$ B into the nucleus.


**Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>15</sub>H<sub>20</sub>O<sub>6</sub>

Batch Molecular Weight: 296.32

Physical Appearance: White solid

**Batch Molecular Structure:**



**Storage:** Store at +4°C

**Solubility & Usage Info:**

ethanol to 30 mM

This product is supplied as a lyophilized solid and may be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

**Stability and Solubility Advice:**

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

**SOLIDS:** Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

**SOLUTIONS:** We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

**References:**

**Krishnaswamy et al** (2010) Lutein protects HT-29 cells against deoxynivalenol-induced oxidative stress and apoptosis: prevention of NF- $\kappa$ B nuclear localization and down regulation of NF- $\kappa$ B and cyclo-oxygenase-2 expression. *Free Radic.Biol.Med.* **49** 50. PMID: 20347963.

**Shi et al** (2009) Role of GRP78/BiP degradation and ER stress in deoxynivalenol-induced interleukin-6 upregulation in the macrophage. *Toxicol.Sci.* **109** 247. PMID: 19336499.

**Bae and Pestka** (2008) Deoxynivalenol induces p38 interaction with the ribosome in monocytes and macrophages. *Toxicol.Sci.* **105** 59. PMID: 18502741.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

**bio-techne.com**

info@bio-techne.com

techsupport@bio-techne.com

**North America**

Tel: (800) 343 7475

**China**

info.cn@bio-techne.com

Tel: +86 (21) 52380373

**Europe Middle East Africa**

Tel: +44 (0)1235 529449

**Rest of World**

www.tocris.com/distributors

Tel:+1 612 379 2956