



# **Certificate of Analysis**

www.tocris.com

Product Name: L-Ascorbic acid Catalog No.: 4055 Batch No.: 2

CAS Number: 50-81-7 EC Number: 200-066-2

IUPAC Name: 3-Oxo-L-gulofuranolactone

## 1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_6H_8O_6$ .Batch Molecular Weight:176.12Physical Appearance:White solid

**Solubility:** water to 500 mM

Storage: Store at RT

**Batch Molecular Structure:** 

#### 2. ANALYTICAL DATA

HPLC: Shows 99.8% purity

<sup>1</sup>H NMR: Consistent with structure Mass Spectrum: Consistent with structure

**Optical Rotation:**  $[\alpha]_D = +20$  (Concentration = 10, Solvent = Water)

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 40.92 4.58 Found 41.11 4.55

www.tocris.com/distributors Tel:+1 612 379 2956

## **Product Information**

Print Date: Oct 11th 2024

www.tocris.com

Product Name: L-Ascorbic acid Catalog No.: 4055 2

CAS Number: 50-81-7 EC Number: 200-066-2

IUPAC Name: 3-Oxo-L-gulofuranolactone

#### **Description:**

L-Ascorbic acid is an inhibitor of  $\text{Ca}_{\text{v}}3.2$  channels (IC $_{50}$  = 6.5  $\mu\text{M}$ ); displays no effect on  $\text{Ca}_{\text{v}}3.1$  or  $\text{Ca}_{\text{v}}3.3$  channels heterologously expressed in HEK 293 cells. Also enhances the generation of induced pluripotent stem cells (iPSCs) from mouse and human somatic cells by increasing reprogramming efficiency. Commonly used antifade reagent in live cell microscopy. Naturally occurring antioxidant. L-Ascorbic acid synthesized to Ancillary Material Grade also available. For more information about how L-Ascorbic acid may be used, see our protocol: Highly Efficient Generation of CiPSCs from MEFs Please see product specific page on www.tocris.com for full description.

## **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>6</sub>H<sub>8</sub>O<sub>6</sub>. Batch Molecular Weight: 176.12 Physical Appearance: White solid

**Minimum Purity:** ≥99%

#### **Batch Molecular Structure:**

о он он

Storage: Store at RT

## Solubility & Usage Info:

water to 500 mM

### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. \*Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

#### References:

**Cordes** *et al* (2011) Mechanisms and advancement of antifading agents for fluorescence microscopy and single-molecule spectroscopy. Phys.Chem.Chem.Phys. *13* 6699. PMID: 21311807.

Esteban et al (2010) Vitamin C enhances the generation of mouse and human induced pluripotent stem cells. Cell Stem Cell 6 71. PMID: 20036631.

**Nelson** *et al* (2007) Molecular mechanisms of subtype-specific inhibition of neuronal T-type calcium channels by ascorbate. J.Neurosci. **27** 12577. PMID: 18003836.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use