



Certificate of Analysis

www.tocris.com

Product Name: DIM Catalog No.: 4502 Batch No.: 1

CAS Number: 1968-05-4

IUPAC Name: 3,3'-Methylenediindole

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: $C_{17}H_{14}N_2$.0.1 H_2 O

Batch Molecular Weight: 248.11
Physical Appearance: Yellow solid

Solubility: DMSO to 100 mM

ethanol to 50 mM

Storage: Store at +4°C

Batch Molecular Structure:

HNNH

2. ANALYTICAL DATA

Melting Point:

HPLC:

Shows 99.5% purity

Consistent with atwest

¹H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 82.3 5.69 11.29 Found 82.14 5.68 11.31



Product Information

Print Date: Jun 15th 2017

www.tocris.com

Product Name: DIM Catalog No.: 4502 Batch No.: 1

CAS Number: 1968-05-4

IUPAC Name: 3,3'-Methylenediindole

Description:

Activator of Chk2 that causes G_2/M cell cycle arrest in various cancer cell lines. Antiproliferative and inducer of apoptosis; promotes proteasomal degradation of Cdc25C and Cdk1. Inhibits phosphorylation of EGFR and downstream activation of ERK.


Physical and Chemical Properties:

Batch Molecular Formula: $C_{17}H_{14}N_2$.0.1 H_2O

Batch Molecular Weight: 248.11 Physical Appearance: Yellow solid

Minimum Purity: >99%

Batch Molecular Structure:



Storage: Store at +4°C

Solubility & Usage Info:

DMSO to 100 mM ethanol to 50 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Kandala et al (2012) Blocking epidermal growth factor receptor activation by 3,3'-diindolylmethane suppresses ovarian tumor growth in vitro and in vivo. J.Pharmacol.Exp.Ther. **341** 24. PMID: 22205686.

Shorey et al (2012) 3,3'-diindolylmethane induces G1 arrest and apoptosis in human acute T-cell lymphoblastic leukemia cells. PLoS One. **7** e34975. PMID: 22514694.

Kandala & Srivastava (2010) Activation of checkpoint kinase 2 by 3,3'-diindolylmethane is required for causing G₂/M cell cycle arrest in human ovarian cancer cells. Mol.Pharmacol. **78** 297. PMID: 20444961.