## **bio-techne**<sup>®</sup> TOCRIS

## **Certificate of Analysis**

#### www.tocris.com

Print Date: Dec 20th 2024

#### Product Name: Cytarabine

147-94-4 CAS Number: IUPAC Name:

4-Amino-1-β-D-arabinofuranosyl-2(1*H*)-pyrimidinone

#### 1. PHYSICAL AND CHEMICAL PROPERTIES

**Batch Molecular Formula: Batch Molecular Weight: Physical Appearance:** Solubility:

 $C_9H_{13}N_3O_5$ . 243.22 White solid

Storage: **Batch Molecular Structure:** 

# $NH_2$ HO

οн


#### 2. ANALYTICAL DATA

HPLC: <sup>1</sup>H NMR: Mass Spectrum: **Optical Rotation: Microanalysis:** 

Shows 99.6% purity Consistent with structure Consistent with structure  $[\alpha]_D$  = +155.8 (Concentration = 0.5, Solvent = Water) Carbon Hydrogen Nitrogen Theoretical 44.45 5.39 17.28 Found 44.36 5.42 17.13

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com	North America	China	Europe Middle East Africa	Rest of World
info@bio-techne.com techsupport@bio-techne.com	Tel: (800) 343 7475	info.cn@bio-techne.com Tel: +86 (21) 52380373	Tel: +44 (0)1235 529449	www.tocris.com/distributors Tel:+1 612 379 2956



Batch No.: 6

Catalog No.: 4520 EC Number: 205-705-9

## biotechne TOCRIS

#### www.tocris.com

Batch No.: 6

#### Product Name: Cytarabine

CAS Number: 147-94-4

IUPAC Name: 4-Amino-1- $\beta$ -D-arabinofuranosyl-2(1*H*)-pyrimidinone

#### **Description:**


Cytarabine is a nucleoside analog of deoxycytidine; inhibits DNA replication by incorporating into DNA ( $IC_{50} = 0.04 \mu$ M in L1210 and CEM cell lines). Displays no inhibitory effects on RNA synthesis. Causes S phase cell cycle arrest in ML-1 cell lines; cytotoxic in L5817Y leukemia cells. Antineoplastic and antileukemic agent.

#### **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>9</sub>H<sub>13</sub>N<sub>3</sub>O<sub>5</sub>. Batch Molecular Weight: 243.22 Physical Appearance: White solid

#### Minimum Purity: ≥98%

#### **Batch Molecular Structure:**



#### **References:**

Sampath et al (2006) Pharmacodynamics of cytar. alone and in combination with 7-hydroxystaurosporine (UCN-01) in AML blasts in vitro and during a clinical trial. Blood. **107** 2517. PMID: 16293603.

**Breistøl** *et al* (1999) Antitumor activity of P-4055 (elaidic acid-cytarabine) compared to cytar. in metastatic and s.c. human tumor xenograft models. Cancer Res. **59** 2944. PMID: 10383159.

Yang et al (1986) Effect of cytar. upon metabolism of phospholipids in leukemia L5178Y cells. Acta Pharmacol. Sin. 7 368. PMID: 2954403.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com	North America	China	Europe Middle East Africa	Rest of World
info@bio-techne.com techsupport@bio-techne.com	Tel: (800) 343 7475	info.cn@bio-techne.com Tel: +86 (21) 52380373	Tel: +44 (0)1235 529449	www.tocris.com/distributors Tel:+1 612 379 2956

### Storage: Store at +4°C

CAUTION - This product is light sensitive and we recommend that the solid material and any solutions obtained are protected from exposure to light.

Catalog No.: 4520

EC Number: 205-705-9

#### Solubility & Usage Info:

water to 100 mM DMSO to 50 mM

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. \*Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.