

Product Name: Resiquimod

Catalog No.: 4536


Batch No.: 3

CAS Number: 144875-48-9

IUPAC Name: 1-[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]-2-methylpropan-2-ol

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₁₇H₂₂N₄O₂
Batch Molecular Weight: 314.38
Physical Appearance: Off White solid
Solubility: DMSO to 100 mM
ethanol to 50 mM
Storage: Store at +4°C
Batch Molecular Structure:



2. ANALYTICAL DATA

HPLC: Shows 99.6% purity
¹H NMR: Consistent with structure
Mass Spectrum: Consistent with structure

Microanalysis:

	Carbon	Hydrogen	Nitrogen
Theoretical	64.95	7.05	17.82
Found	65.08	7.07	17.84

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

Product Name: Resiquimod

Catalog No.: 4536

Batch No.: 3

CAS Number: 144875-48-9

IUPAC Name: 1-[4-Amino-2-(ethoxymethyl)-1*H*-imidazo[4,5-*c*]quinolin-1-yl]-2-methylpropan-2-ol

Description:

Toll-like receptor 7 (TLR7) agonist. Induces upregulation of IL-6, IL-12, IFN- γ and iNOS expression in mouse bone marrow-derived macrophages (BMMs). Inhibits RANKL-induced osteoclast differentiation in mouse BMMs and human peripheral blood monocytes. Also promotes the differentiation of MDSCs into macrophages and dendritic cells. Proinflammatory and antiviral.

Physical and Chemical Properties:


Batch Molecular Formula: C₁₇H₂₂N₄O₂

Batch Molecular Weight: 314.38

Physical Appearance: Off White solid

Minimum Purity: >98%

Batch Molecular Structure:



Storage: Store at +4°C

Solubility & Usage Info:

DMSO to 100 mM
ethanol to 50 mM

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

References:

Lee *et al* (2014) Resiquimod, a TLR7/8 agonist, promotes differentiation of myeloid-derived suppressor cells into macrophages and dendritic cells. *Arch.Pharm.Res.* **37** 1234. PMID: 24748512.

Miyamoto *et al* (2012) R848, a toll-like receptor 7 agonist, inhibits osteoclast differentiation but not survival or bone-resorbing function of mature osteoclasts. *Cytotechnology.* **64** 331. PMID: 22358541.

Dockrell and Kinghorn (2001) Imiquimod and resiquimod as novel immunomodulators. *J.Antimicrob.Chemother.* **48** 751. PMID: 11733457.

Wagner *et al* (1999) Modulation of TH1 and Th2 cytokine production with the immune response modifiers, R-848 and imiquimod. *Cell Immunol.* **191** 10. PMID: 9918682.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com
techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com
Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors
Tel:+1 612 379 2956