

Product Name: FSLLRY-NH2

Catalog No.: 4751

Batch No.: 5

CAS Number: 245329-02-6

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: C₃₉H₆₀N₁₀O₈
Batch Molecular Weight: 796.97
Physical Appearance: White lyophilised solid
Net Peptide Content: 78%
Counter Ion: TFA
Solubility: Soluble to 1 mg/ml in water
Storage: Store at -20°C
Peptide Sequence: Phe-Ser-Leu-Leu-Arg-Tyr-NH₂

2. ANALYTICAL DATA

HPLC: Shows 97.6% purity
Mass Spectrum: Consistent with structure

3. AMINO ACID ANALYSIS DATA

Amino Acid Theoretical		Actual		Amino Acid Theoretical		Actual	
Ala				Lys			
Arg	1.00	0.98	Met				
Asx			Phe	1.00	1.00		
Cys			Pro				
Glx			Ser	1.00	0.78		
Gly			Thr				
His			Trp				
Ile			Tyr	1.00	0.83		
Leu	2.00	2.02	Val				

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel: +1 612 379 2956

Product Name: FSLLRY-NH2**Catalog No.:** 4751**Batch No.:** 5

CAS Number: 245329-02-6

Description:

Selective PAR₂ peptide antagonist. Reverses taxol-induced mechanical allodynia, heat hyperalgesia and PKC activation in ICR mice. Blocks ERK activation and collagen production in isolated cardiac fibroblasts. Also reduces symptoms in a mouse model of dermatophyte-associated itch.

Physical and Chemical Properties:Batch Molecular Formula: C₃₉H₆₀N₁₀O₈

Batch Molecular Weight: 796.97

Physical Appearance: White lyophilised solid

Peptide Sequence:Phe-Ser-Leu-Leu-Arg-Tyr-NH₂**Storage:** Store at -20°C**Solubility & Usage Info:**

Soluble to 1 mg/ml in water

This product is supplied as a lyophilized solid and may be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Net Peptide Content: 78% (Remaining weight made up of counterions and residual water).**Counter Ion:** TFA**Stability and Solubility Advice:**

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such as Cys, Met, Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 µm filter to remove potential bacterial contamination whenever possible.

References:

Andoh et al (2012) Involvement of serine protease and proteinase-activated receptor 2 in dermatophyte-associated itch in mice. *J.Pharmacol.Exp.Ther.* **343** 91. PMID: 22761302.

Chen et al (2011) Proteinase-activated receptor 2 sensitizes transient receptor potential vanilloid 1, transient receptor potential vanilloid 4, and transient receptor potential ankyrin 1 in PacT.-induced neuropathic pain. *Neuroscience* **193** 440. PMID: 21763756.

McLarty et al (2011) Tryptase/protease-activated receptor 2 interactions induce selective mitogen-activated protein kinase signaling and collagen synthesis by cardiac fibroblasts. *Hypertension* **58** 264. PMID: 21730297.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com

info@bio-techne.com

techsupport@bio-techne.com

North America

Tel: (800) 343 7475

China

info.cn@bio-techne.com

Tel: +86 (21) 52380373

Europe Middle East Africa

Tel: +44 (0)1235 529449

Rest of World

www.tocris.com/distributors

Tel:+1 612 379 2956