

# **Certificate of Analysis**

Print Date: Jan 14th 2016

www.tocris.com

Catalog No.: 5277 Batch No.: 1

Product Name: CCG 203971 CAS Number: 1443437-74-8

IUPAC Name: N-(4-Chlorophenyl)-1-[3-(2-furanyl)benzoyl]-3-piperidinecarboxamide

#### 1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula:  $C_{23}H_{21}CIN_2O_3$ 

Batch Molecular Weight: 408.88

Physical Appearance: White solid

Solubility: DMSO to 50 mM

Storage: Store at +4°C

**Batch Molecular Structure:** 

#### 2. ANALYTICAL DATA

HPLC: Shows 98.3% purity

<sup>1</sup>H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

Microanalysis:

Carbon Hydrogen Nitrogen

Theoretical 67.56 5.18 6.85 Found 67.26 5.1 6.8



## **Product Information**

Print Date: Jan 14<sup>th</sup> 2016

www.tocris.com

Product Name: CCG 203971 Catalog No.: 5277 Batch No.: 1

CAS Number: 1443437-74-8

IUPAC Name: N-(4-Chlorophenyl)-1-[3-(2-furanyl)benzoyl]-3-piperidinecarboxamide

#### **Description:**

Antifibrotic agent; inhibits fibrosis by targeting the MRTF/SRF gene transcription pathway. Selectively inhibits proliferation of SSc-derived dermal fibroblasts but not that of normal fibroblasts. Inhibits expression of CTGF,  $\alpha\text{-SMA}$ , and COL1A2 in SSc fibroblasts as well as in LPA and in TGF $\beta$ -stimulated fibroblasts. Prevents bleomycin-induced skin thickening and collagen deposition in vivo. Also suppresses PC-3 cell migration in scratch wound assays (IC $_{50}$  = 4.2  $\mu\text{M}$ ).

#### **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>23</sub>H<sub>21</sub>CIN<sub>2</sub>O<sub>3</sub> Batch Molecular Weight: 408.88

Physical Appearance: White solid

### Batch Molecular Structure:

Minimum Purity: >98%

NH CI Storage: Store at +4°C

#### Solubility & Usage Info:

DMSO to 50 mM

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

#### References:

**Bell** et al (2013) Optimization of novel nipecotic bis(amide) inhibitors of the Rho/MKL1/SRF transcriptional pathway as potential antimetastasis agents. Bioorg.Med.Chem.Lett. **23** 3826. PMID: 23707258.

**Haak** *et al* (2014) Targeting the myofibroblast genetic switch: inhibitors of myocardin-related transcription factor/serum response factor-regulated gene transcription prevent fibrosis in a murine model of skin injury. J.Pharmacol.Exp.Ther. **349** 480. PMID: 24706986.

**Michel** *et al* (2014) Hope for disease-modifying treatment of systemic sclerosis/scleroderma. J.Pharmacol.Exp.Ther. *350* 480. PMID: 25053235.