biotechne[®] TOCRIS

Certificate of Analysis

www.tocris.com

Print Date: Jan 3rd 2025

Catalog No.: 5941 Batch No.: 8

Product Name: OD1

1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: Batch Molecular Weight: Physical Appearance: Counter Ion: Solubility: Storage: Peptide Sequence: C₃₀₈H₄₆₆N₉₀O₉₅S₈ 7206.1 White solid TFA Soluble to 1 mg/ml in water Store at -20°C Gly-Val-Arg-Asp-Ala-Tyr-IIe-Ala-Asp-Asp-

Lys-Asn-Cys-Val-Tyr-Thr-Cys-Ala-Ser-Asn-

Gly-Tyr-Cys-Asn-Thr-Glu-Cys-Thr-Lys-Asn-

Gly-Ala-Glu-Ser-Gly-Tyr-Cys-Gln-Trp-Ile-

Gly-Arg-Tyr-Gly-Asn-Ala-Cys-Trp-Cys-Ile-

Lys-Leu-Pro-Asp-Glu-Val-Pro-Ile-Arg-Ile-

Pro-Gly-Lys-Cys-Arg-NH2

2. ANALYTICAL DATA

HPLC: Mass Spectrum: Shows 98.2 % purity Consistent with structure

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.comNorth AmericaChinaEurope Middle East AfricaRest of Worldinfo@bio-techne.comTel: (800) 343 7475info.cn@bio-techne.comTel: +44 (0)1235 529449www.tocris.com/distributorstechsupport@bio-techne.comTel: +86 (21) 52380373Tel: +44 (0)1235 529449tel: +1612 379 2956

biotechne[®] TOCRIS

www.tocris.com

Print Date: Jan 3rd 2025

Product Name: OD1

Description:

OD1 is a potent rat Na_v1.7, human Na_v1.4 and rat Na_v1.6 channel activator (EC₅₀ values are 7, 10 and 47 nM, respectively). Exhibits minimal activation at mammalian Na_v1.2, Na_v1.3 and Na_v1.5 (EC₅₀ values >3 μ M). Inhibits fast inactivation on all channels. Increases peak currents at all voltages and stimulates a persistent Na⁺ current at hNa_v1.7 channel. Increases hyperpolarization at Na_v1.4 and Na_v1.6 channels. Induces spontaneous pain in vivo.

Physical and Chemical Properties:

Batch Molecular Formula: C₃₀₈H₄₆₆N₉₀O₉₅S₈ Batch Molecular Weight: 7206.1 Physical Appearance: White solid

Peptide Sequence:

Gly-Val-Arg-Asp-Ala-Tyr-Ile-Ala-Asp-Asp-

Lys-Asn-Cys-Val-Tyr-Thr-Cys-Ala-Ser-Asn-

Gly-Tyr-Cys-Asn-Thr-Glu-Cys-Thr-Lys-Asn-

Gly-Ala-Glu-Ser-Gly-Tyr-Cys-Gln-Trp-lle-

Gly-Arg-Tyr-Gly-Asn-Ala-Cys-Trp-Cys-Ile-

Lys-Leu-Pro-Asp-Glu-Val-Pro-Ile-Arg-Ile-

Pro-Gly-Lys-Cys-Arg-NH₂

References:

Durek *et al* (2013) Chemical engineering and structural and pharmacological characterization of the α -scorpion toxin OD1. ACS Chem.Biol. **8** 1215. PMID: 23527544.

Maertens *et al* (2006) Potent modulation of the voltage-gated sodium channel Na_v1.7 by OD1, a toxin from the scorpion *Odonthobuthus doriae*. Mol.Pharmacol. **70** 405. PMID: 16641312.

Jalali *et al* (2005) OD1, the first toxin isolated from the venom of the scorpion Odonthobuthus doriae active on voltage-gated Na⁺ channels. FEBS Lett. **579** 4181. PMID: 16038905.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com	North America	China	Europe Middle East Africa	Rest of World
info@bio-techne.com techsupport@bio-techne.com	Tel: (800) 343 7475	info.cn@bio-techne.com Tel: +86 (21) 52380373	Tel: +44 (0)1235 529449	www.tocris.com/distributors Tel:+1 612 379 2956

Catalog No.: 5941 Batch No.: 8

Storage: Store at -20°C

Solubility & Usage Info:

Soluble to 1 mg/ml in water

This product is supplied in lyophilized form. It may appear as a solid, gel or film and be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Counter Ion: TFA

Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such Cys, Met,Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2 μ m filter to remove potential bacterial contamination whenever possible.