## biotechne<sup>®</sup> TOCRIS

## **Certificate of Analysis**

# Print Date: Sep 3<sup>rd</sup> 2024

www.tocris.com

Product Name: OD1

Catalog No.: 5941 Batch No.: 9

### 1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: Batch Molecular Weight: Physical Appearance: Counter Ion: Solubility: Storage: Peptide Sequence: C<sub>308</sub>H<sub>466</sub>N<sub>90</sub>O<sub>95</sub>S<sub>8</sub> 7206.1 White solid TFA Soluble to 1 mg/ml in water Store at -20°C Gly-Val-Arg-Asp-Ala-Tyr-IIe-Ala-Asp-Asp-

Lys-Asn-Cys-Val-Tyr-Thr-Cys-Ala-Ser-Asn-

Gly-Tyr-Cys-Asn-Thr-Glu-Cys-Thr-Lys-Asn-

Gly-Ala-Glu-Ser-Gly-Tyr-Cys-Gln-Trp-Ile-

Gly-Arg-Tyr-Gly-Asn-Ala-Cys-Trp-Cys-Ile-

Lys-Leu-Pro-Asp-Glu-Val-Pro-Ile-Arg-Ile-

Pro-Gly-Lys-Cys-Arg-NH2

### 2. ANALYTICAL DATA

HPLC: Mass Spectrum: Shows 98.4% purity Consistent with structure

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

## biotechne<sup>®</sup> TOCRIS

### www.tocris.com

#### Product Name: OD1

#### **Description:**

OD1 is a potent rat Na<sub>v</sub>1.7, human Na<sub>v</sub>1.4 and rat Na<sub>v</sub>1.6 channel activator (EC<sub>50</sub> values are 7, 10 and 47 nM, respectively). Exhibits minimal activation at mammalian Na<sub>v</sub>1.2, Na<sub>v</sub>1.3 and Na<sub>v</sub>1.5 (EC<sub>50</sub> values >3  $\mu$ M). Inhibits fast inactivation on all channels. Increases peak currents at all voltages and stimulates a persistent Na<sup>+</sup> current at hNa<sub>v</sub>1.7 channel. Increases hyperpolarization at Na<sub>v</sub>1.4 and Na<sub>v</sub>1.6 channels. Induces spontaneous pain in vivo.

#### **Physical and Chemical Properties:**

Batch Molecular Formula:  $C_{308}H_{466}N_{90}O_{95}S_8$ Batch Molecular Weight: 7206.1 Physical Appearance: White solid

#### **Peptide Sequence:**

Gly-Val-Arg-Asp-Ala-Tyr-Ile-Ala-Asp-Asp-

Lys-Asn-Cys-Val-Tyr-Thr-Cys-Ala-Ser-Asn-

Gly-Tyr-Cys-Asn-Thr-Glu-Cys-Thr-Lys-Asn-

Gly-Ala-Glu-Ser-Gly-Tyr-Cys-Gln-Trp-lle-

Gly-Arg-Tyr-Gly-Asn-Ala-Cys-Trp-Cys-Ile-

Lys-Leu-Pro-Asp-Glu-Val-Pro-Ile-Arg-Ile-

Pro-Gly-Lys-Cys-Arg-NH<sub>2</sub>

#### **References:**

**Durek** *et al* (2013) Chemical engineering and structural and pharmacological characterization of the α-scorpion toxin OD1. ACS Chem.Biol. **8** 1215. PMID: 23527544.

**Maertens** *et al* (2006) Potent modulation of the voltage-gated sodium channel Na<sub>v</sub>1.7 by OD1, a toxin from the scorpion *Odonthobuthus doriae*. Mol.Pharmacol. **70** 405. PMID: 16641312.

Jalali *et al* (2005) OD1, the first toxin isolated from the venom of the scorpion Odonthobuthus doriae active on voltage-gated Na<sup>+</sup> channels. FEBS Lett. **579** 4181. PMID: 16038905.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

| bio-techne.com                                    | North America       | China  | Europe Middle East Africa | Rest of World                                      |
|---|---------------------|--|---------------------------|--|
| info@bio-techne.com<br>techsupport@bio-techne.com | Tel: (800) 343 7475 | info.cn@bio-techne.com<br>Tel: +86 (21) 52380373 | Tel: +44 (0)1235 529449   | www.tocris.com/distributors<br>Tel:+1 612 379 2956 |

#### Catalog No.: 5941

9

Storage: Store at -20°C

#### Solubility & Usage Info:

Soluble to 1 mg/ml in water

This product is supplied in lyophilized form. It may appear as a solid, gel or film and be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

#### Counter Ion: TFA

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such Cys, Met,Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2  $\mu$ m filter to remove potential bacterial contamination whenever possible.