



# **Certificate of Analysis**

www.tocris.com

Product Name: Caerulein Catalog No.: 6264 Batch No.: 9

CAS Number: 17650-98-5

## 1. PHYSICAL AND CHEMICAL PROPERTIES

**Batch Molecular Formula:**  $C_{58}H_{73}N_{13}O_{21}S_2$ 

Batch Molecular Weight: 1352.4

**Physical Appearance:** White lyophilised solid **Counter Ion:** Ammonium Acetate

**Solubility:** Soluble to 1 mg/ml in water

Storage: Store at -20°C

Peptide Sequence: Glp-Gln-Asp-Tyr(SO<sub>3</sub>H)-Thr-Gly-Trp-Met-Asp-

Phe-NH<sub>2</sub>

2. ANALYTICAL DATA

HPLC: Shows 97.0% purity

Mass Spectrum: Consistent with structure

3. AMINO ACID ANALYSIS DATA

Amino Acid	l Theoretica	l Actual	Amino Acid	Theoretica	Actual
Ala			Lys		
Arg			Met	1.00	0.97
Asx	2.00	2.03	Phe	1.00	0.99
Cys			Pro		
Glx	2.00	2.03	Ser		
Gly	1.00	0.97	Thr	1.00	0.86
His			Trp	1.00	0.62
lle			Tyr	1.00	1.01
Leu			Val		

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use



# **Product Information**

Print Date: Mar 12th 2024

www.tocris.com

Product Name: Caerulein Catalog No.: 6264 9

CAS Number: 17650-98-5

#### **Description:**

Caerulein is a CCK agonist. Stimulates biosynthesis of pancreatic enzymes in AR4-2J pancreatic cells in vitro. Induces acute pancreatitis in mice. Also enhances proteasome activity in Gdeg mice.

### **Physical and Chemical Properties:**

Batch Molecular Formula:  $C_{58}H_{73}N_{13}O_{21}S_2$ 

Batch Molecular Weight: 1352.4

Physical Appearance: White lyophilised solid

#### **Peptide Sequence:**

Glp-Gln-Asp-Tyr(SO<sub>3</sub>H)-Thr-Gly-Trp-Met-Asp-Phe-NH<sub>2</sub> Storage: Store at -20°C

# Solubility & Usage Info:

Soluble to 1 mg/ml in water

This product is supplied in lyophilized form. It may appear as a solid, gel or film and be very hard to visualize. Solutions should be made by adding solvent directly to the vial. The vial should then be vortexed vigorously to ensure the product has completely dissolved.

Counter Ion: Ammonium Acetate

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Peptides in solution are much less stable than in lyophilized form. This is especially true for peptides whose sequences contain amino acids such Cys, Met,Trp, Asn, Gln, and N-terminal Glu.

Therefore we recommend storing peptides in solution for as short a time as possible. Avoid repeated freeze thaw cycles by dividing the peptide solution into aliquots and storing the aliquots at -20°C. Any portion of an aliquot unused after thawing should be discarded.

Peptides stored in solution can occasionally be susceptible to bacterial degradation. We recommend using sterile solutions or passing the peptide solution through a 0.2  $\mu$ m filter to remove potential bacterial contamination whenever possible.

#### References:

**Murakami** et al (2017) Yes-associated protein mediates immune reprogramming in pancreatic ductal adenocarcinoma. Oncogene **36** 1232. PMID: 27546622.

Furuyama et al (2016) Proteasome activity is required for the initiation of precancerous pancreatic lesions. Sci.Rep. 6 27044. PMID: 27244456.

**Pradel** et al (1993) Caerulein and gastrin(2-17 ds) regulate differently synthesis of secretory enzymes, mRNA levels and cell proliferation in pancreatic acinar cells (AR4-2J). Biochem.J. **290** 219. PMID: 7679894.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use