# biotechne<sup>®</sup> TOCRIS

# Print Date: May 29<sup>th</sup> 2024

Catalog No.: 7122

# **Certificate of Analysis**

### www.tocris.com

Batch No.: 3

### Product Name: Thioflavin T

CAS Number: 2390-54-7

IUPAC Name: 2-[4-(Dimethylamino)phenyl]-3,6-dimethylbenzothiazolium chloride

## 1. PHYSICAL AND CHEMICAL PROPERTIES

Batch Molecular Formula: Batch Molecular Weight: Physical Appearance: Solubility:

**Batch Molecular Structure:** 

C<sub>17</sub>H<sub>19</sub>CIN<sub>2</sub>S 318.86 Yellow solid DMSO to 5 mM water to 10 mM Store at -20°C

CF

### 2. ANALYTICAL DATA

Storage:

HPLC: <sup>1</sup>H NMR: Mass Spectrum: Net product content:

Shows 97.8% purity Consistent with structure Consistent with structure 88%

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.comNorth AmericaChinaEurope Middle East AfricaRest of Worldinfo@bio-techne.comTel: (800) 343 7475info.cn@bio-techne.comTel: +44 (0)1235 529449www.tocris.com/distributorstechsupport@bio-techne.comTel: +86 (21) 52380373Tel: +44 (0)1235 529449tel: +1 612 379 2956

# biotechne<sup>®</sup> TOCRIS

### www.tocris.com

3

### Product Name: Thioflavin T

CAS Number: 2390-54-7

IUPAC Name: 2-[4-(Dimethylamino)phenyl]-3,6-dimethylbenzothiazolium chloride

#### **Description:**


Key information: Thioflavin T is a cell-permeable fluorescent amyloid stain. Used for: staining of insoluble senile plaques of A $\beta$  in brain tissues, monitoring the kinetics of in vitro polyglutamine amyloid formation of tNhtt-42Q aggregates in a Huntington's disease cell model. Thioflavin T is used to confirm formation of  $\beta$  sheet structure from mutant huntingtin exon-1 aggregates (Ex1Q48) in vitro. Application: fluorescence microscopy. Properties and Photophysical Data: Thioflavin T is a cationic benzothiazole. It increases in fluorescence upon strong binding (K<sub>i</sub> = 580 nM) to the stacked  $\beta$  sheets of amyloid fibrils. Excitati... Please see product specific page on www.tocris.com for full description.

#### **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>17</sub>H<sub>19</sub>ClN<sub>2</sub>S Batch Molecular Weight: 318.86 Physical Appearance: Yellow solid

#### Minimum Purity: ≥95%

#### **Batch Molecular Structure:**



#### References:

# Schindler (2021) Small, seeding-competent huntingtin fibrils are prominent aggregate species in brains of zQ175 Huntington's disease knock-in mice. Front.Neurosci. **15** 682172. PMID: 34239412.

Lee *et al* (2019) Fluorescence chemicals to detect insoluble and soluble amyloid-β aggregates. ACS Chem.Neurosci. **10** 2647. PMID: 31009195.

**Doi** *et al* (2008) RNA-binding protein TLS is a major nuclear aggregate-interacting protein in huntingtin exon 1 with expanded polyglutamine-expressing cells. J.Biol.Chem. **283** 6489. PMID: 18167354.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com	North America	China	Europe Middle East Africa	Rest of World
info@bio-techne.com techsupport@bio-techne.com	Tel: (800) 343 7475	info.cn@bio-techne.com Tel: +86 (21) 52380373	Tel: +44 (0)1235 529449	www.tocris.com/distributors Tel:+1 612 379 2956

CAUTION - This product is light sensitive and we recommend that the solid material and any solutions obtained are protected from exposure to light.

Catalog No.: 7122

#### Solubility & Usage Info:

Storage: Store at -20°C

DMSO to 5 mM water to 10 mM

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. \*Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.