

# **Certificate of Analysis**

Print Date: Apr 7th 2022

www.tocris.com

Product Name: CSN5i-3-NEG Catalog No.: 7532 Batch No.: 1

isopropyl-1H-pyrazole-5-carboxamide

#### 1. PHYSICAL AND CHEMICAL PROPERTIES

**Batch Molecular Formula:**  $C_{28}H_{29}F_2N_5O_2.\frac{1}{2}H_2O$ 

**Batch Molecular Weight:** 514.58

Physical Appearance: White solid

Solubility: DMSO to 100 mM Storage: Store at -20°C

**Batch Molecular Structure:** 

## 2. ANALYTICAL DATA

HPLC: Shows 99.7% purity
Chiral HPLC: Shows 100% purity

<sup>1</sup>H NMR: Consistent with structure

Mass Spectrum: Consistent with structure

Microanalysis: Carbon Hydrogen Nitrogen

Theoretical 65.36 5.88 13.61 Found 65.11 5.68 13.41

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use



# **Product Information**

Print Date: Apr 7<sup>th</sup> 2022

www.tocris.com

Product Name: CSN5i-3-NEG Catalog No.: 7532 Batch No.: 1

isopropyl-1*H*-pyrazole-5-carboxamide

#### **Description:**

CSN5i-3-NEG is the negative control for CSN5i-3 (Cat. No.7089).

### **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>28</sub>H<sub>29</sub>F<sub>2</sub>N<sub>5</sub>O<sub>2</sub>.½H<sub>2</sub>O

Batch Molecular Weight: 514.58 Physical Appearance: White solid

Minimum Purity: ≥98%

#### **Batch Molecular Structure:**

Storage: Store at -20°C

### Solubility & Usage Info:

DMSO to 100 mM

#### Stability and Solubility Advice:

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. Our standard recommendations are:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

#### References:

Majolée et al (2019) CSN5 inhibition triggers inflammatory signaling and Rho/ROCK-dependent loss of endothelial integrity. Sci.Rep. 9 8131. PMID: 31148579.

**Zhang** *et al* (2017) COPS5 inhibition arrests the proliferation and growth of serous ovarian cancer cells via the elevation of p27 level. Biochem.Biophys.Res.Commun. *493* 85. PMID: 28919423.

Schlierf et al (2016) Targeted inhibition of the COP9 signalosome for treatment of cancer. Nat Commun. 7 13166. PMID: 27774986.