# **biotechne**<sup>®</sup> TOCRIS

### Print Date: Feb 21st 2024

### **Certificate of Analysis**

### www.tocris.com

#### Product Name: FAI

CAS Number: 1415238-77-5 **IUPAC Name:** 1H-Imidazol-1-yl(2-methyl-3-furanyl)methanone

### 1. PHYSICAL AND CHEMICAL PROPERTIES

**Batch Molecular Formula: Batch Molecular Weight: Physical Appearance:** Solubility: Storage: **Batch Molecular Structure:**   $C_9H_8N_2O_2$ 176.17 White crystalline solid DMSO to 100 mM Store at -20°C

### 2. ANALYTICAL DATA

<sup>1</sup>H NMR: Mass Spectrum: Consistent with structure Consistent with structure

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.com **North America** China **Europe Middle East Africa Rest of World** info@bio-techne.com Tel: (800) 343 7475 info.cn@bio-techne.com Tel: +44 (0)1235 529449 www.tocris.com/distributors Tel: +86 (21) 52380373 techsupport@bio-techne.com Tel:+1 612 379 2956



Catalog No.: 8011 Batch No.: 1

## biotechne TOCRIS

### Product Name: FAI

Catalog No.: 8011

1

Print Date: Feb 21st 2024

www.tocris.com

CAS Number:1415238-77-5IUPAC Name:1H-Imidazol-1-yl(2-methyl-3-furanyl)methanone

### **Description:**

FAI is a reagent for RNA SHAPE-MaP experiments in live cells. Can be used to map RNA structures in vivo by forming stable 2'-OH adducts with RNA, blocking reverse transcriptase elongation. In vitro, FAI activity can be quenched with Dithiothreitol (Cat. No. 3154) and can be used to probe the secondary structure of mouse embryonic stem cell 5S rRNA. At elevated temperatures, FAI forms more stable adducts and exhibits less hydrolysis than NAI (Cat. No 7004).

### **Physical and Chemical Properties:**

Batch Molecular Formula: C<sub>9</sub>H<sub>8</sub>N<sub>2</sub>O<sub>2</sub> Batch Molecular Weight: 176.17 Physical Appearance: White crystalline solid

**Batch Molecular Structure:** 



Storage: Store at -20°C

Solubility & Usage Info: DMSO to 100 mM

#### **Stability and Solubility Advice:**

Some solutions can be difficult to obtain and can be encouraged by rapid stirring, sonication or gentle warming (in a 45-60°C water bath).

Information concerning product stability, particularly in solution, has rarely been reported and in most cases we can only offer a general guide. \*Unless contradicted by product-specific protocols or instructions, our standard recommendations apply:

SOLIDS: Provided storage is as stated on the product label and the vial is kept tightly sealed, the product can be stored for up to 6 months from date of receipt.

SOLUTIONS: We recommend that stock solutions, once prepared, are stored aliquoted in tightly sealed vials at -20°C or below and used within 1 month. Wherever possible solutions should be made up and used on the same day.

### **References:**

**Chan** *et al* (2017) Comparative analysis reveals furoyl in vivo selective hydroxyl acylation analyzed by primer extension reagents form stable ribosyl ester adducts. Biochemistry **56** 18111. PMID: 28319368.

Spitale et al (2013) RNA SHAPE analysis in living cells. Nat.Chem.Biol. 9 18. PMID: 23178934.

Caution - Not Fully Tested • Research Use Only • Not For Human or Veterinary Use

bio-techne.comNorth AmericaChinaEurope Middle East AfricaRest of Worldinfo@bio-techne.comTel: (800) 343 7475info.cn@bio-techne.comTel: +44 (0)1235 529449www.tocris.com/distributorstechsupport@bio-techne.comTel: +86 (21) 52380373Tel: +44 (0)1235 529449tel: +1 612 379 2956