

Product Datasheet

EPCR Antibody (M2) - Azide and BSA Free H00010544-M03

Unit Size: 0.1 mg

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/H00010544-M03

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/H00010544-M03



H00010544-M03

EPCR Antibody (M2) - Azide and BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	M2
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	IgG purified
Buffer	In 1x PBS, pH 7.4


Product Description	
Description	Novus Biologicals Mouse EPCR Antibody (M2) - Azide and BSA Free (H00010544-M03) is a monoclonal antibody validated for use in IHC, WB, ELISA and ICC/IF. Anti-EPCR Antibody: Cited in 7 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	10544
Gene Symbol	PROCR
Species	Human, Mouse
Reactivity Notes	Other species not tested.
Specificity/Sensitivity	PROCR - protein C receptor, endothelial (EPCR)
Immunogen	PROCR (AAH14451.1, 1 a.a. ~ 238 a.a) full-length recombinant protein with GST tag. MW of the GST tag alone is 26 KDa. MLTLLPILLLSGWAFCSQDASDGLQRLHMLQISYFRDPYHVWYQGNASLGGH LTHVLEGPDTNTTIIQLQPLQEPESWARTQSGLQSYLLQFHGLVRLVHQERTLA FPLTIRCFLGCELPPEGSRAHVFFEVAVNGSSSFVSFRPERALWQADTQVTSGV VTFTLQQLNAYNRTRYELREFLEDTCVQYVQKHISAENTKGSQTSRSYTSLVLG VLVGGFIIAGVAVGIFLCTGGRRRC
Notes	This product is produced by and distributed for Abnova, a company based in Taiwan.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Sandwich ELISA
Recommended Dilutions	Western Blot 1:500, ELISA 1:100-1:2000, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence 1:10-1:500, Immunohistochemistry-Paraffin 1:10-1:500, Sandwich ELISA
Application Notes	Antibody reactive against recombinant protein for Western Blot. Has also been used for immunofluorescence, immunohistochemistry (paraffin), and ELISA.




Images


Immunocytochemistry/Immunofluorescence: EPCR Antibody (M2) [H00010544-M03] - Analysis of monoclonal antibody to PROCR on HeLa cell. Antibody concentration 10 ug/ml



Immunohistochemistry-Paraffin: EPCR Antibody (M2) [H00010544-M03] - Analysis of monoclonal antibody to PROCR on formalin-fixed paraffin-embedded human malignant lymphoma, diffuse large B tissue. Antibody concentration 3 ug/ml



Sandwich ELISA: EPCR Antibody (M2) [H00010544-M03] - Detection limit for recombinant GST tagged PROCR is 3 ng/ml as a capture antibody.



Publications

Ku SK, Zhou W, Lee W et al. Anti-inflammatory effects of hyperoside in human endothelial cells and in mice. *Inflammation* 2014-08-06 [PMID: 25097077]

Lee W, Ku SK, Bae JS. Emodin-6-O-b-D-glucoside down-regulates endothelial protein C receptor shedding. *Arch Pharm Res.* 2013-04-16 [PMID: 23588681]

Ku SK, Han MS, Bae JS. Down-regulation of endothelial protein C receptor shedding by persicarin and isorhamnetin-3-O-galactoside. *Thromb Res.* 2013-05-30 [PMID: 23726966]

Ku SK, Yang EJ, Song KS, Bae JS. Rosmarinic acid down-regulates endothelial protein C receptor shedding in vitro and in vivo. *Food Chem Toxicol.* 2013-06-14 [PMID: 23774263]

Ku SK, Lee W, Yoo H et al. Inhibitory effects of epi-sesamin on endothelial protein C receptor shedding in vitro and in vivo. *Inflamm Res.* 2013-07-25 [PMID: 23884236]

Ku SK, Han MS, Bae JS. Sulforaphane inhibits endothelial protein C receptor shedding in vitro and in vivo. *Vascul Pharmacol.* 2014-07-09 [PMID: 25016099]

Kwak S, Han MS, Bae JS. Aspalathin and nothofagin from rooibos (*Aspalathus linearis*) inhibit endothelial protein C receptor shedding in vitro and in vivo. *Fitoterapia* 2014-12-12 [PMID: 25510322]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to H00010544-M03

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/H00010544-M03

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

