

# Product Datasheet

## APE Antibody (13B8E5C2) - BSA Free NB100-116

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 3 Publications: 111**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-116](http://www.novusbio.com/NB100-116)

Updated 9/9/2025 v.20.1

**Earn rewards for product  
reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-116](http://www.novusbio.com/reviews/destination/NB100-116)



**NB100-116**

APE Antibody (13B8E5C2) - BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	13B8E5C2
Preservative	0.02% Sodium Azide
Isotype	IgG2b
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	37 kDa

Product Description	
Description	Novus Biologicals Mouse APE Antibody (13B8E5C2) - BSA Free (NB100-116) is a monoclonal antibody validated for use in IHC, WB, ELISA, ICC/IF, Simple Western, IP and ChIP. Anti-APE Antibody: Cited in 106 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	328
Gene Symbol	APEX1
Species	Human, Mouse, Rat, Primate
Immunogen	Purified human APE1 [Uniprot: P27695]

Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, ELISA, Gel Super Shift Assays, Immunoblotting, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation, Proximity Ligation Assay, Chromatin Immunoprecipitation (ChIP), Knockdown Validated
Recommended Dilutions	Western Blot 1:100-1:2000, Simple Western 1:25, ELISA reported in scientific literature (PMID 21769563), Immunohistochemistry 1:100, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunoprecipitation 1:10 - 1:500. Use reported in scientific literature (PMID 35286386), Immunohistochemistry-Paraffin 1:100, Immunohistochemistry-Frozen 1:10-1:500, Immunoblotting reported in scientific literature (PMID 27608656), Gel Super Shift Assays reported in scientific literature, Proximity Ligation Assay reported in scientific literature (PMID 27808278), Chromatin Immunoprecipitation (ChIP) 1:10-1:500, Knockdown Validated




**Application Notes**

In Western blot, this antibody detects a single band at 37 kDa. In IHC, it can be competitively inhibited from recognizing the APE1 antigen in tissues using APE1 protein. It can also be used on frozen and fixed-paraffin sections and cytospin preps. In IHC-P, staining was observed in the nucleus of a human breast cancer xenograft. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.


See [Simple Western Antibody Database](#) for Simple Western validation: Tested in HeLa lysate 0.1 mg/mL, separated by Size, antibody dilution of 1:25. Separated by Size-Wes, Sally Sue/Peggy Sue.

**Images**

Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - Whole cell protein from human HeLa, A431, mouse Neuro2A and rat PC12 cells was separated on a 12% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2.0 ug/mL anti-APE-1 in block buffer and detected with an anti-mouse HRP secondary antibody using chemiluminescence.




Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - Repression of p21 by APE1 in p53-null cells and effect of ectopic p53 in this repression. Representative Western analysis of p53, APE1, p21 and alpha-Tubulin levels in the same HCT116p53null cells. Image collected and cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0068467>) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - HeLa cells stained NB100-116 (Green) detected with DyLight Fluor 488 conjugated anti-rabbit IgG secondary antibody. Nuclei are counterstained with Hoechst 33258 (Blue).



**Immunohistochemistry: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - Immunohistochemical staining of APE1.** A. Nuclear staining in the luminal epithelium of normal breast ducts and lobules. B. Low, and C. high nuclear APE1 expression in invasive breast cancer. (magnification x400). Image collected and cropped by CiteAb from the following publication ([//dx.plos.org/10.1371/journal.pone.0099528](https://dx.plos.org/10.1371/journal.pone.0099528)) licensed under a CC-BY license.




**Chromatin Immunoprecipitation (ChIP): APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - Association of p53 and APE1 on p53-binding sites in p21 promoter.** Re-ChIP analysis (first IP with alpha-APE1 and the second IP with alpha-p53 antibody) showing simultaneous recruitment of APE1 and p53 in control vs. EPE treated cells; \*: p value <0.05 (n=2) calculated based on APE1/p53 enriched DNA from control vs. etoposide treated cells. Image collected and cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0068467>) licensed under a CC-BY license.




**Simple Western: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - Image shows a specific band for APE1 in 0.1 mg/mL of HeLa lysate.** This experiment was performed under reducing conditions using the 12-230kDa separation system. \* Non-specific interaction with the 230 kDa Simple Western standard may be seen with this antibody.



**Immunocytochemistry/Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - HeLa cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.05% Triton X-100.** The cells were incubated with anti-APE (13B8E5C2) at 5 ug/mL overnight at 4C and detected with an anti-mouse DyLight 488 (Green) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - Ovarian Cancer cell lines.




Immunocytochemistry/Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - Immunocytochemical detection of APE-ref-1 in breast cancer cell line MDA MB 231.




Immunohistochemistry-Paraffin: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE Antibody (13B8E5C2) [NB100-116] - APE1 antibody was tested in human breast cancer xenograft using DAB with hematoxylin counterstain.




Chromatin Immunoprecipitation: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Association of APE1 & AP4 in p21 proximal promoter region. (A & B) ChIP Real Time PCR analysis showing relative enrichment of (A) APE1 & (B) AP4 in p21 proximal promoter containing AP4-responsive E-Box elements in HCT116p53null cells. (C) Re-ChIP (first IP with  $\alpha$ -APE1 & the second IP with  $\alpha$ -AP4 antibody) analysis showing simultaneous recruitment of APE1 & AP4 in this promoter region. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/23874636>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Effect of APE1 on p21 activation in p53 WT cells. (A) Western analysis for APE1 level in WT APE1 &  $\Delta 42$  APE1 overexpressing cells. (B) Real Time RT-PCR analysis showing relative quantitation of p21 transcript level in etoposide-treated WT APE1-overexpressing HCT116WT cells &  $\Delta 42$  APE1 overexpressing cells; \*: p value <0.05 (n=2) calculated from control (empty vector transfection) vs. WT APE1 overexpression, WT APE1 overexpression along with etoposide treatment or  $\Delta 42$  overexpression. (C) Luciferase activity in cells co-transfected with empty or WT APE1-expression vector & p21 promoter-luciferase construct; \*: p value <0.05 (n=2) calculated from control (empty vector transfection) vs. APE1 overexpression. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/23874636>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Analysis of single strand break repair in MCF-7 monolayer & mammosphere cell populations. (a) Cells were exposed to 4 Gy 60Co  $\gamma$ -radiation & the relative degree of single-strand breakage (SSB) was determined by alkaline single-cell gel electrophoresis (comet assay) immediately after exposure & at the times indicated after exposure. (b) The 'comets' (n of about 100) were categorized according to the NIH LISTSERV (Comet Assay Interest Group web site) in which type 1 comets display the least DNA damage & type 5 the most. The error bars represent the mean  $\pm$  standard error of the mean in both panels. The comets of the unirradiated cells are labeled Cont. (c) Expression of proteins involved in SSB repair in response to ionizing radiation. Lysates were prepared from unirradiated cells & from cells harvested one hour after exposure to 1 or 10-Gy 60Co  $\gamma$ -radiation & analyzed by immunoblotting with antibodies against several SSB repair proteins.  $\alpha$ -Actin served as a loading control. Image collected & cropped by CiteAb from the following publication (<http://breast-cancer-research.biomedcentral.com/articles/10.1186/bcr2583>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Intracellular APE1/Ref-1 was targeted to the plasma membrane in response to acetylation. (A & B) Whole cell lysates were immunoprecipitated using an anti-acetyl lysine antibody, followed by immunoblotting with the polyclonal anti-APE1/Ref-1 antibody. The blots were stripped & re-probed with anti- $\beta$ -actin & APE1/Ref-1 antibody to ensure equal protein loading. Similar results were obtained from replicate experiments. Column, mean (n = 3); bars, SE. \*, p < 0.05, significantly different compared with control or between group by one-way ANOVA followed by Bonferroni's multiple comparison test. (C,D) Membrane fractions or whole cell lysates were prepared from the TSA-treated cells. Immunoblotting for APE1/Ref-1 was performed using the polyclonal anti-APE1/Ref-1 antibody. Blots were stripped & re-probed with anti-N-cadherin & anti- $\beta$ -actin antibodies to control for differences in protein loading. Fold changes in the levels of APE1/Ref-1 in the plasma membrane fraction relative to the control are shown. †, indicates molecular marker (N-cadherin) of left image. Column mean (n = 3); bars, SE. \*, p < 0.05 indicates a significantly different result compared with control or between groups by one-way ANOVA followed by Bonferroni's multiple comparison test. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31261750>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - AcAPE1 is exclusively associated w/ chromatin & remains bound to condensed chromosomes. (A & B) Asynchronous normal lung fibroblast IMR90 cells & lung adenocarcinoma A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs, counterstained w/ DAPI, & visualized by confocal microscopy & 3D SIM. (C) Colocalization of AcAPE1 w/ histone H3 or active enhancer-specific histone marker acetylated H3K27 (H3K27Ac). (D) BJ-hTERT cells serum starved for 72 h & then fixed at different time points. Cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs & counterstained w/ anti-TO-PRO-3 iodide Ab. (E) Mitotic A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 & visualized by 3D SIM. (F) BJ-hTERT cells either serum starved for 72 h (G0/G1 phase), treated w/ nocodazole (mitotic cells) or aphidicolin (G1/S phase synchronized cells), or untreated, & whole-cell extracts isolated using 150 mM or 300 mM salt-containing lysis buffer. Western blot analysis for anti-APE1 & anti-AcAPE1 levels performed. Anti-HSC70 used as loading control. (G) A proximal ligation assay performed w/ mouse anti-APE1 & rabbit anti-APE1 (mAPE1 & Rabbit-APE1), mouse anti-mouse APE1 & rabbit anti-AcAPE1 (mAPE1 & rAcAPE1), & rabbit anti-AcAPE1 & mouse anti-histone H3 (mHistone H3 & rAcAPE1) to confirm the chromatin association of AcAPE1. Mouse IgG (mIgG) & rabbit anti-AcAPE1 used as a control. At least 50 cells counted for PLA foci. (H) Colocalization of p300 & AcAPE1 on chromatin (DAPI). (I) HCT116 cells transfected w/ E1A & mutant E1A, & at 48 h after transfection, IF performed. Cells immunostained w/ anti-p300 & anti-APE1 or anti-AcAPE1 & counterstained w/ DAPI. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - XPF is epistatic to APE in the gemcitabine resistance. The expression of XPF and/or APE was suppressed by siRNAs in HeLa cells & the cellular sensitivity to gemcitabine was examined. XPF- or APE-suppressed HeLa cells (closed square & closed triangle, respectively) showed sensitivity to gemcitabine. Cosuppression of XPF & APE (closed circle with dashed line) resulted in the sensitivity to gemcitabine similar to the sensitivity induced by the suppression of XPF or APE individually. A control siRNA (siControl) was used as a control (open diamond). Three independent experiments were performed & averages of surviving fraction are plotted. The error bars show standard deviations. The differences in the gemcitabine sensitivity between the control cells & the cells treated with siXPF, siAPE, or siXPF+siAPE are statistically significant at 10 nM & 50 nM with  $p < 0.05$ . The western blots showed a significant suppression of XPF (more than 95%) & ~75% reduction in the expression of APE with the siRNA treatment. The cosuppression experiments with two siRNAs, siXPF & siAPE, resulted in similar levels of suppression of each protein induced by individual siRNA (more than 95% reduction in XPF & ~85% reduction in APE). Tubulin was used as a protein loading control. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30941207>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Plasma membrane associated APE1/Ref-1 is bound to ABCA1 in response to acetylation. Cells transiently expressing wild type APE1/Ref-1-FLAG or mutant APE1/Ref-1(K6/7R)-FLAG were treated with 1  $\mu$ M TSA for 1 h. (A) Whole cell lysates were immunoprecipitated using the monoclonal anti-ABCA1 antibody, followed by immunoblot with the anti-FLAG antibody. (B) For reverse immunoprecipitation, cell lysates were immunoprecipitated with anti-APE1/Ref-1 antibody followed by immunoblot analysis with the polyclonal anti-ABCA1 antibody. Blots were stripped & re-probed with anti-ABCA1 or FLAG antibodies to ensure equal protein loading & no contamination of cellular proteins. Similar results were observed in replicate experiments. Columns, mean (n = 2-3); bars, SE. \*, p < 0.05 indicates a significantly different result from control cells according to unpaired t-tests. (C) The binding between APE1/Ref-1 & ABCA1 in the plasma membrane was visualized using with a Duolink II PLA system with primary polyclonal anti-APE1/Ref-1 & monoclonal anti-ABCA1 antibodies (PLA<sup>+</sup>). The PLA-specific fluorescence which represents the APE1/Ref-1-ABCA1 signal, & the DAPI nuclear staining are in red & blue, respectively. The experiment was repeated multiple times with similar results; the data shown here are from a representative experiment. Optical slices were examined using a 40 $\times$  oil immersion objective with a 2 $\times$  zoom factor. Scale bar, 20  $\mu$ m ( $\times$ 80). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31261750>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE1 acetylation enhances its stability on chromatin & its interaction with downstream BER proteins. (A) Colocalization of ligase III & AcAPE1 in A549 cells. Cells were immunostained with anti-ligase III & anti-AcAPE1 Abs. (B) WT or K5R mutant APE1-overexpressing HEK293T cells were treated with TSA-nicotinamide (NAM) for 6 h or not treated, & the nuclear extract was immunoprecipitated using anti-FLAG Ab & immunoblotted with anti-XRCC1 & anti-FLAG Abs. (C) A549 cells were fixed with paraformaldehyde before (top) or after treatment with Triton X-100 (0.5%) (middle) or Triton X-100 plus salt (100 mM KCl) (bottom) & immunostained with anti-APE1 or anti-AcAPE1 Abs & counterstained with DAPI. (D) Acetylation of APE1 induces a conformational change in APE1. The distinct intrinsic fluorescence emission spectra of APE1 & AcAPE1 at 280 nm are shown. A.U., absorbance units. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




**Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - AcAPE1 is exclusively associated w/ chromatin & remains bound to condensed chromosomes. (A & B) Asynchronous normal lung fibroblast IMR90 cells & lung adenocarcinoma A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs, counterstained w/ DAPI, & visualized by confocal microscopy & 3D SIM. (C) Colocalization of AcAPE1 w/ histone H3 or active enhancer-specific histone marker acetylated H3K27 (H3K27Ac). (D) BJ-hTERT cells serum starved for 72 h & then fixed at different time points. Cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs & counterstained w/ anti-TO-PRO-3 iodide Ab. (E) Mitotic A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 & visualized by 3D SIM. (F) BJ-hTERT cells either serum starved for 72 h (G0/G1 phase), treated w/ nocodazole (mitotic cells) or aphidicolin (G1/S phase synchronized cells), or untreated, & whole-cell extracts isolated using 150 mM or 300 mM salt-containing lysis buffer. Western blot analysis for anti-APE1 & anti-AcAPE1 levels performed. Anti-HSC70 used as loading control. (G) A proximal ligation assay performed w/ mouse anti-APE1 & rabbit anti-APE1 (mAPE1 & Rabbit-APE1), mouse anti-mouse APE1 & rabbit anti-AcAPE1 (mAPE1 & rAcAPE1), & rabbit anti-AcAPE1 & mouse anti-histone H3 (mHistone H3 & rAcAPE1) to confirm the chromatin association of AcAPE1. Mouse IgG (mIgG) & rabbit anti-AcAPE1 used as a control. At least 50 cells counted for PLA foci. (H) Colocalization of p300 & AcAPE1 on chromatin (DAPI). (I) HCT116 cells transfected w/ E1A & mutant E1A, & at 48 h after transfection, IF performed. Cells immunostained w/ anti-p300 & anti-APE1 or anti-AcAPE1 & counterstained w/ DAPI. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.**




**Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE1 is acetylated after binding to AP sites in the chromatin. (A) BJ-hTERT cells were treated with MX (50 mM) for the indicated times. IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. (B) HCT116 cells were treated with various doses of MX for 30 min, IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. (C) HCT116 cells were treated with 50 mM MX for 30 min, IF was performed using anti-OGG1, & counterstaining with DAPI was used. (D) BJ-hTERT cells pretreated with 50 mM MX for 30 min or not pretreated were exposed to MMS (2 mM) for 1 h. IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. Confocal microscopy was used to visualize the AcAPE1 levels in control cells & cells treated with MMS or MX, or both. (E) ChIP with anti-OGG1 antibody followed by Western blotting (ChIP-on-Western) was performed to examine the association of AcAPE1 & ligase III on chromatin after induction of DNA damage with GO. (F) The association of AcAPE1 with the endogenous p21 promoter in control or MMS- or MX-treated cells was examined by promoter-directed ChIP using anti-AcAPE1. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.**




**Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116]** - APE1 is acetylated after binding to AP sites in the chromatin. (A) BJ-hTERT cells were treated with MX (50 mM) for the indicated times. IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. (B) HCT116 cells were treated with various doses of MX for 30 min, IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. (C) HCT116 cells were treated with 50 mM MX for 30 min, IF was performed using anti-OGG1, & counterstaining with DAPI was used. (D) BJ-hTERT cells pretreated with 50 mM MX for 30 min or not pretreated were exposed to MMS (2 mM) for 1 h. IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. Confocal microscopy was used to visualize the AcAPE1 levels in control cells & cells treated with MMS or MX, or both. (E) ChIP with anti-OGG1 antibody followed by Western blotting (ChIP-on-Western) was performed to examine the association of AcAPE1 & ligase III on chromatin after induction of DNA damage with GO. (F) The association of AcAPE1 with the endogenous p21 promoter in control or MMS- or MX-treated cells was examined by promoter-directed ChIP using anti-AcAPE1. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




**Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116]** - AcAPE1 is exclusively associated w/ chromatin & remains bound to condensed chromosomes. (A & B) Asynchronous normal lung fibroblast IMR90 cells & lung adenocarcinoma A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs, counterstained w/ DAPI, & visualized by confocal microscopy & 3D SIM. (C) Colocalization of AcAPE1 w/ histone H3 or active enhancer-specific histone marker acetylated H3K27 (H3K27Ac). (D) BJ-hTERT cells serum starved for 72 h & then fixed at different time points. Cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs & counterstained w/ anti-TO-PRO-3 iodide Ab. (E) Mitotic A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 & visualized by 3D SIM. (F) BJ-hTERT cells either serum starved for 72 h (G0/G1 phase), treated w/ nocodazole (mitotic cells) or aphidicolin (G1/S phase synchronized cells), or untreated, & whole-cell extracts isolated using 150 mM or 300 mM salt-containing lysis buffer. Western blot analysis for anti-APE1 & anti-AcAPE1 levels performed. Anti-HSC70 used as loading control. (G) A proximal ligation assay performed w/ mouse anti-APE1 & rabbit anti-APE1 (mAPE1 & Rabbit-APE1), mouse anti-mouse APE1 & rabbit anti-AcAPE1 (mAPE1 & rAcAPE1), & rabbit anti-AcAPE1 & mouse anti-histone H3 (mHistone H3 & rAcAPE1) to confirm the chromatin association of AcAPE1. Mouse IgG (mIgG) & rabbit anti-AcAPE1 used as a control. At least 50 cells counted for PLA foci. (H) Colocalization of p300 & AcAPE1 on chromatin (DAPI). (I) HCT116 cells transfected w/ E1A & mutant E1A, & at 48 h after transfection, IF performed. Cells immunostained w/ anti-p300 & anti-APE1 or anti-AcAPE1 & counterstained w/ DAPI. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - AcAPE1 is exclusively associated w/ chromatin & remains bound to condensed chromosomes. (A & B) Asynchronous normal lung fibroblast IMR90 cells & lung adenocarcinoma A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs, counterstained w/ DAPI, & visualized by confocal microscopy & 3D SIM. (C) Colocalization of AcAPE1 w/ histone H3 or active enhancer-specific histone marker acetylated H3K27 (H3K27Ac). (D) BJ-hTERT cells serum starved for 72 h & then fixed at different time points. Cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs & counterstained w/ anti-TO-PRO-3 iodide Ab. (E) Mitotic A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 & visualized by 3D SIM. (F) BJ-hTERT cells either serum starved for 72 h (G0/G1 phase), treated w/ nocodazole (mitotic cells) or aphidicolin (G1/S phase synchronized cells), or untreated, & whole-cell extracts isolated using 150 mM or 300 mM salt-containing lysis buffer. Western blot analysis for anti-APE1 & anti-AcAPE1 levels performed. Anti-HSC70 used as loading control. (G) A proximal ligation assay performed w/ mouse anti-APE1 & rabbit anti-APE1 (mAPE1 & Rabbit-APE1), mouse anti-mouse APE1 & rabbit anti-AcAPE1 (mAPE1 & rAcAPE1), & rabbit anti-AcAPE1 & mouse anti-histone H3 (mHistone H3 & rAcAPE1) to confirm the chromatin association of AcAPE1. Mouse IgG (mIgG) & rabbit anti-AcAPE1 used as a control. At least 50 cells counted for PLA foci. (H) Colocalization of p300 & AcAPE1 on chromatin (DAPI). (I) HCT116 cells transfected w/ E1A & mutant E1A, & at 48 h after transfection, IF performed. Cells immunostained w/ anti-p300 & anti-APE1 or anti-AcAPE1 & counterstained w/ DAPI. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Repression of p21 by APE1 in p53-null cells & effect of ectopic p53 in this repression. (A & B) Real Time RT-PCR analysis showing relative quantitation of p21 transcript level in (A) HCT116p53null cells with WT &  $\Delta$ 42 APE1 overexpression; \*: p value (n=4) calculated from control (empty vector transfection) vs. WT or  $\Delta$ 42 APE1 overexpression, & (B) control (control siRNA) vs. APE1-depleted HCT116p53null cells; \*: p value <0.05 (n=4) calculated from control vs. APE1-depleted cells. (C) Effect of ectopic p53 expression on p21 transcript level in control vs. APE1-depleted HCT116p53null cells. First, cells were transfected with control siRNA or APE1 siRNA, the next day both the cell types were again transfected with empty vector or p53 expression vector & after 48 hrs the cells were harvested; signal from empty vector transfection in both control & APE1-depleted cells were set as reference samples; \*: p value <0.05 (n=3) calculated based on the effect of ectopic p53 expression over empty vector transfection in control vs. APE1-depleted cells. (D) Effect of APE1 depletion in control (empty vector transfected) vs. ectopic p53-expressing HCT116p53null cells; the same experiment was performed as in C but analyzed differently; signal from control siRNA-transfected cells in both empty vector transfected & ectopic p53 expressing cases were set as reference samples; \*: p value <0.05 (n=3) calculated based on the effect of APE1-depletion in empty vector transfected vs. ectopic p53 expressing cells. (E) Representative Western analysis of p53, APE1, p21 &  $\alpha$ -Tubulin levels in the same HCT116p53null cells as in B–D. (F & G) Real Time RT-PCR analysis of p21 level in Saos2 cells as in C & D. \*: p value <0.05 (n=2). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/23874636>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - APE1 is acetylated after binding to AP sites in the chromatin. (A) BJ-hTERT cells were treated with MX (50 mM) for the indicated times. IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. (B) HCT116 cells were treated with various doses of MX for 30 min, IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. (C) HCT116 cells were treated with 50 mM MX for 30 min, IF was performed using anti-OGG1, & counterstaining with DAPI was used. (D) BJ-hTERT cells pretreated with 50 mM MX for 30 min or not pretreated were exposed to MMS (2 mM) for 1 h. IF was performed using anti-APE1 & anti-AcAPE1, & counterstaining with DAPI was used. Confocal microscopy was used to visualize the AcAPE1 levels in control cells & cells treated with MMS or MX, or both. (E) ChIP with anti-OGG1 antibody followed by Western blotting (ChIP-on-Western) was performed to examine the association of AcAPE1 & ligase III on chromatin after induction of DNA damage with GO. (F) The association of AcAPE1 with the endogenous p21 promoter in control or MMS- or MX-treated cells was examined by promoter-directed ChIP using anti-AcAPE1. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Intracellular APE1/Ref-1 was targeted to the plasma membrane in response to acetylation. (A & B) Whole cell lysates were immunoprecipitated using an anti-acetyl lysine antibody, followed by immunoblotting with the polyclonal anti-APE1/Ref-1 antibody. The blots were stripped & re-probed with anti- $\beta$ -actin & APE1/Ref-1 antibody to ensure equal protein loading. Similar results were obtained from replicate experiments. Column, mean (n = 3); bars, SE. \*, p < 0.05, significantly different compared with control or between group by one-way ANOVA followed by Bonferroni's multiple comparison test. (C,D) Membrane fractions or whole cell lysates were prepared from the TSA-treated cells. Immunoblotting for APE1/Ref-1 was performed using the polyclonal anti-APE1/Ref-1 antibody. Blots were stripped & re-probed with anti-N-cadherin & anti- $\beta$ -actin antibodies to control for differences in protein loading. Fold changes in the levels of APE1/Ref-1 in the plasma membrane fraction relative to the control are shown. †, indicates molecular marker (N-cadherin) of left image. Column mean (n = 3); bars, SE. \*, p < 0.05 indicates a significantly different result compared with control or between groups by one-way ANOVA followed by Bonferroni's multiple comparison test. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31261750>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




**Immunocytochemistry/ Immunofluorescence: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - AcAPE1 is exclusively associated w/ chromatin & remains bound to condensed chromosomes. (A & B)** Asynchronous normal lung fibroblast IMR90 cells & lung adenocarcinoma A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs, counterstained w/ DAPI, & visualized by confocal microscopy & 3D SIM. (C) Colocalization of AcAPE1 w/ histone H3 or active enhancer-specific histone marker acetylated H3K27 (H3K27Ac). (D) BJ-hTERT cells serum starved for 72 h & then fixed at different time points. Cells immunostained w/ anti-APE1 & anti-AcAPE1 Abs & counterstained w/ anti-TO-PRO-3 iodide Ab. (E) Mitotic A549 cells immunostained w/ anti-APE1 & anti-AcAPE1 & visualized by 3D SIM. (F) BJ-hTERT cells either serum starved for 72 h (G0/G1 phase), treated w/ nocodazole (mitotic cells) or aphidicolin (G1/S phase synchronized cells), or untreated, & whole-cell extracts isolated using 150 mM or 300 mM salt-containing lysis buffer. Western blot analysis for anti-APE1 & anti-AcAPE1 levels performed. Anti-HSC70 used as loading control. (G) A proximal ligation assay performed w/ mouse anti-APE1 & rabbit anti-APE1 (mAPE1 & Rabbit-APE1), mouse anti-mouse APE1 & rabbit anti-AcAPE1 (mAPE1 & rAcAPE1), & rabbit anti-AcAPE1 & mouse anti-histone H3 (mHistone H3 & rAcAPE1) to confirm the chromatin association of AcAPE1. Mouse IgG (mIgG) & rabbit anti-AcAPE1 used as a control. At least 50 cells counted for PLA foci. (H) Colocalization of p300 & AcAPE1 on chromatin (DAPI). (I) HCT116 cells transfected w/ E1A & mutant E1A, & at 48 h after transfection, IF performed. Cells immunostained w/ anti-p300 & anti-APE1 or anti-AcAPE1 & counterstained w/ DAPI. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27994014>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



**Chromatin Immunoprecipitation: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Association of p53 & APE1 on p53-binding sites in p21 promoter. (A) p21 promoter structure showing p53 & AP4 binding sites. (B) ChIP Real Time PCR analysis showing relative enrichment ( $2^{-\Delta\Delta CT}$ ) of APE1-immunoprecipitated DNA over that from control IgG in p21 promoter regions containing p53 binding sites 1 & 2 in HCT116WT cells. (C) Re-ChIP analysis (first IP with  $\alpha$ -APE1 & the second IP with  $\alpha$ -p53 antibody) showing simultaneous recruitment of APE1 & p53 in control vs. etoposide treated cells; \*: p value <0.05 (n=2) calculated based on APE1/p53 enriched DNA from control vs. etoposide treated cells. (D) Western analysis of FLAG immunoprecipitate (IP) to detect APE1-associated p53 & FLAG (APE1) from empty vector vs. FLAG-tagged WT APE1 or FLAG-tagged N $\Delta$ 33 APE1 transfected HCT116WT cells (left panel) & from control vs. etoposide-treated WT APE1-FLAG transfected cells (right panel). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/23874636>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.**



Western Blot: APE Antibody (13B8E5C2) - BSA Free [NB100-116] - Gemcitabine-induced recruitment of XPF to chromatin depends on APE. HCT116, APE-suppressed HCT116 (HCT116 shAPE), & XPF-deficient HCT116 (HCT116 g4-10) cells were treated with 1  $\mu$ M of gemcitabine & chromatin fractions were isolated in the indicated time. The presence of XPF & APE was detected by western blots. The asterisk (\*) shows a cross-reacted protein with the anti-XPF antibody. Histone H2AX was used as a loading control. Gemcitabine-induced recruitment of XPF to chromatin (lanes 1-3) was compromised in HCT116 shAPE cell line (lanes 4-5). The chromatin-bound APE is not changed by the gemcitabine treatment (lanes 4-6, Supplementary Figure 5). A signal of XPF (and APE) was normalized with a signal of H2AX in each chromatin fraction using ImageJ software. Then a ratio of chromatin-bound XPF with gemcitabine to XPF in control was determined & depicted as bar graphs. Three independent experiments were performed & averages of the ratio at indicated time points were plotted. The error bars show standard deviations. Only the difference in the chromatin-bound XPF between chromatin from control experiments & chromatin that was incubated one hour with 1  $\mu$ M gemcitabine is statistically significant in HCT116 ( $p < 0.05$ ). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30941207>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Muniyandi A, Hartman G, Sishtla K et al. Ref-1 is overexpressed in neovascular eye disease and targetable with a novel inhibitor *Angiogenesis* 2025-04-24 [PMID: 39756006]

Kiran, S;Wireman, RS;Peil, J;Mitchell, DK;Potchanant, ES;Rai, R;Vilseck, JZ;Cao, S;Haiaty, S;Georgiadis, MM;Fishel, ML;Kelley, MR; The absence of Peroxiredoxin-1 in human pancreatic ductal adenocarcinoma (PDAC) markedly reduces cell survival and tumor growth when coupled with the inhibition of Ref-1 redox signaling *Redox biology* 2025-09-02 [PMID: 40902319]

Musiani D, Yücel H, Vallette M et al. Uracil processing by SMUG1 in the absence of UNG triggers homologous recombination and selectively kills BRCA1/2-deficient tumors. *Molecular cell* 2025-02-19 [PMID: 40010343]

Gabriella DH, Anbukkarasi M, Kamakshi S et al. Ref-1 redox activity regulates retinal neovascularization by modulating transcriptional activation of HIF-1 $\alpha$  *FASEB J.* 2025-02-15 [PMID: 39902892] (Immunohistochemistry, Western Blot, Human, Mouse)

Hesbon Z. Amenya, Chiharu Tohyama, Seiichiroh Ohsako Dioxin induces Ahr-dependent robust DNA demethylation of the Cyp1a1 promoter via Tdg in the mouse liver *Scientific Reports* 2016-10-07 [PMID: 27713569]

Shrabasti Roychoudhury, Suravi Pramanik, Hannah L. Harris, Mason Tarpley, Aniruddha Sarkar, Gaele Spagnol, Paul L. Sorgen, Dipanjan Chowdhury, Vimla Band, David Klinkebiel, Kishor K. Bhakat Endogenous oxidized DNA bases and APE1 regulate the formation of G-quadruplex structures in the genome *Proceedings of the National Academy of Sciences of the United States of America* 2020-05-26 [PMID: 32404420]

Chen YH, Kuo YY, You YQ et al. Endonuclease VIII-like 1 deficiency potentiates nigrostriatal dopaminergic neuron degeneration in a male mouse model of Parkinson's disease *Journal of neurochemistry* 2023-02-24 [PMID: 36840377]

Champion J Targeting the REF1/STAT3 axis to treat tuberous sclerosis Thesis 2023-01-01 (Western Blot, Human)

Details:

1:1000 WB dilution

Ito M, Ducasa GM, Molina JD et al. ABCA1 deficiency contributes to podocyte pyroptosis priming via the APE1/IRF1 axis in diabetic kidney disease *Scientific reports* 2023-06-14 [PMID: 37316538] (WB, Mouse)

Details:

1:500 dilution, fresh sections used

Chieffi Baccari G, Falvo S, Di Fiore MM et al. High-fat diet affects autophagy and mitochondrial compartment in rat Harderian gland *Journal of experimental zoology. Part A, Ecological and integrative physiology* 2022-08-04 [PMID: 35927786]

Latino D, Chieffi Baccari G, Di Fiore MM et al. Autophagy and mitochondrial damage in the testis of high-fat diet fed rats *General and comparative endocrinology* 2022-08-13 [PMID: 35973585] (WB, Rat)

Details:

Dilution used for WB 1:1500

Rios-Covian D, Butcher LD, Ablack AL et al. A novel hypomorphic Apex1 mouse model implicates apurinic-apyrimidinic endonuclease 1 in oxidative DNA damage repair in gastric epithelial cells *Antioxidants & redox signaling* 2022-06-25 [PMID: 35754343] (WB)

More publications at <http://www.novusbio.com/NB100-116>

## Procedures

### Western Blot Protocol for APE1 Antibody (NB100-116)

#### Western Blot

1. Gels, Whatman paper, and membranes are soaked in electroblotting buffer (25 mM Tris-HCl; 193 mM glycine; 20% methanol) for 15 minutes prior to transferring
2. Proteins separated on SDS-polyacrylamide gels are transferred onto 0.22 micron nitrocellulose sheets by electroblotting in a Transblot BioRad transfer apparatus in 25 mM Tris, 192 mM Glycine, 20% Methanol at 150 mA (70 V). The transfer is carried out for 1 hour at 4 degrees C.
3. Following protein transfer, the filter is blocked with Blotto [1X TBST (10X TBST = 1.5 M NaCl; 100 mM Tris-HCl, pH 8.0; 0.5% Tween 20; 2% NP-40; 0.2% SDS); 5% Carnation dried milk; 0.02% sodium azide] for 1 hour at room temperature on a rotator.
4. Dilute NB 100-116 (anti-APE/ref-1) in Blotto and incubate with the filter, at 4 degrees C overnight, on a rotator.
5. Wash filter 3 times in 1X TBST (50 mM Tris-HCl, 150 mM NaCl, 0.1% Tween 20, pH 7.5) for 10 minutes at 4 degrees C. Secondary antibody (peroxidase conjugated anti-mouse) is incubated with the blot for 30 minutes at room temperature. Cross-reacting proteins are detected using the Chemiluminescence Western Blotting Kit from Boehringer-Mannheim.

#### NOTE:

HeLa whole cell extracts (NB800-PC1) were used as a positive control for this antibody.

Immunohistochemistry/Immunocytochemistry

The description that follows is for cultured cells but can be used for cytopins.

1. Split cells into 3.5 cm culture dishes for growth.
2. Wash cells with 5 ml PBS.
3. Fix cells with approx. 3 ml Histochoice (Amresco) for 30 min (Cryostat tissues for 45 min) or use 10% formalin for 30 minutes.
4. Rinse cells with 5 ml TBS, wipe plate dry leaving a small circle of buffer and cells. Mark with red pencil.
5. Pre-block the cells for 30 min. with 10% goat serum in TBS (200 ul).
6. Aspirate blocking solution and add NB 100-116 (anti-APE/ref-1), dilution in 10% goat serum.
7. Incubate in humidified chamber for 3 hours (overnight for tissue at 4 degrees C).
8. Incubate the cells with 1:100 diluted secondary antibody (anti-mouse IgG made in goat) in 10% goat serum and TBS for 1 hour in humidified chamber.
9. Wash 2 times with 5 ml TBS for 5 min each.
10. Block with ABC solution for 30 min.
11. Wash 2 times with 5 ml TBS for 5 min each.
12. Incubate with DAB solution until signal develops. Place into dH<sub>2</sub>O. Add coverslip with Aqua-mount. TBS: 50 mM Tris, 150 mM NaCl, pH 7.5, ABC and DAB solutions: Vector laboratories



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-116**

---

NB800-PC1	HeLa Whole Cell Lysate
NBP1-49581	APE1 Redox Inhibitor
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP2-27231	Mouse IgG2b Isotype Control (MPC-11)

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-116](http://www.novusbio.com/reviews/submit/NB100-116)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)



