

Product Datasheet

ANP32B Antibody NB100-1199

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1199

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1199



NB100-1199

ANP32B Antibody


Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Description	Novus Biologicals Goat ANP32B Antibody (NB100-1199) is a polyclonal antibody validated for use in IHC, WB, ELISA and ICC/IF. Anti-ANP32B Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	10541
Gene Symbol	ANP32B
Species	Human, Mouse, Rat, Porcine
Reactivity Notes	Rat reactivity reported in scientific literature (PMID: 25034417).
Specificity/Sensitivity	This antibody does NOT recognize the TNF family member also known as APRIL - Gene ID number 8741. However this antibody may cross-react with ANP32A (GeneID 8125).
Immunogen	Peptide with sequence C-KRKRETDDEGEDD corresponding to C-Terminus according to NP_006392.1.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Peptide ELISA, Knockdown Validated
Recommended Dilutions	Western Blot 0.1 - 0.3 ug/ml, Immunohistochemistry 3 - 5 ug/ml, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 3-5 ug/ml, Peptide ELISA Detection limit 1:16000, Knockdown Validated
Application Notes	WB: Approx. 26 kDa band observed in human tonsil lysates (calculated MW of 28.8 kDa band according to NP_006392.1). IHC-P: Human breast shows nuclear staining in lobular cells.

Images


Western Blot: ANP32B Antibody [NB100-1199] - 0.1 ug/ml) staining of Human Tonsil (A) and Rat (B) and Pig (C) Spleen lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.




Immunocytochemistry/Immunofluorescence: ANP32B Antibody [NB100-1199] - LPS/IFN gamma-stimulated neuron-glia co-cultures (Glia/PAL31+ neuron) showing expression of both iNOS and PAL31. Image collected and cropped by CiteAb from the following publication (jbiomedsci.com/content/21/1/60), licensed under a CC-BY license.




Immunohistochemistry: ANP32B Antibody [NB100-1199] - (C) Representative micrograph showing colocalization of upregulated PAL31 (green) with microglia/macrophage (ED1 positive cells) (D) Representative micrograph showing colocalization of upregulated PAL31 with astroglia (GFAP positive cells). (E) High power image showing the colocalization of upregulated PAL31 with ED1 positive macrophage/microglia (arrowhead) (F) High power image demonstrating the colocalization of upregulated PAL31 with GFAP positive astroglia (arrow). Magnification: 200X (A-D) and 1000X (E&F). Image collected and cropped by CiteAb from the following publication (jbiomedsci.com/content/21/1/60), licensed under a CC-BY license.




Western Blot: ANP32B Antibody [NB100-1199] - (0.1 ug/ml) staining of NIH3T3 cell lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.




Immunohistochemistry-Paraffin: ANP32B Antibody [NB100-1199] - Staining of paraffin embedded Human Breast. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.




Western Blot: ANP32B Antibody [NB100-1199] - Western blot analysis of pal31 siRNA-treated C6 showing knockdown of PAL31 expression by PAL31 silencer using 41.5 (lane1), 83 (lane2), and 166 (lane 3) picomole of pal31 siRNA or 332 picomole negative control. Actin works as a loading control. Image collected and cropped by CiteAb from the following publication (jbiomedsci.com/content/21/1/60), licensed under a CC-BY license.




Immunocytochemistry/ Immunofluorescence: ANP32B Antibody [NB100-1199] - Upregulated PAL31 expression was observed not only in microglia/ macrophage but also in glial cells near the injury site. (A) Normal thoracic spinal cord section showing few ED-1 positive cells (red). (B) Normal thoracic spinal cord section showing GFAP positive cells (red). (C) Representative micrograph showing colocalization of upregulated PAL31 (green) with microglia/macrophage (ED1 positive cells) (D) Representative micrograph showing colocalization of upregulated PAL31 with astroglia (GFAP positive cells). (E) High power image showing the colocalization of upregulated PAL31 with ED1 positive macrophage/microglia (arrowhead) (F) High power image demonstrating the colocalization of upregulated PAL31 with GFAP positive astroglia (arrow). Magnification: 200X (A-D) & 1000X (E&F). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/25034417>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: ANP32B Antibody [NB100-1199] - Overexpression of PAL31 in C6 did not affect the cell proliferative activity. (A) GFP/bright field-superimposed micrograph in GFP overexpressing cells showing > 50% transfective efficiency (B) GFP-PAL31/bright field-micrograph in PAL31 overexpressing C6 showing correct nuclear localization of GFP-tagged PAL31 & transfection efficiency. (C) Western blot analysis showing the expression of GFP (~27 kDa), GFP-tagged PAL31 (~58 kDa), PCNA (~34 kDa), endogenous PAL31 (~31 kDa) & Actin (~43 kDa) in GFP- or PAL31-C6 cultures. The level of PCNA, a protein expressing in nuclear during DNA synthesis, in cells did not alter after overexpression of GFP or GFP-tagged PAL31. Actin works as a loading control. (D) MTT assay in the transfected cells at 4 different time intervals (from Ctrl to 72 hours). Ctrl represents the cells after subculture & overnight incubation. The data in each time points were analyzed by two-way ANOVA & Bonferroni post hoc test. No significance, compared GFP & GFP tagged PAL31 groups. n = 3. Magnification 100X (A-B). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/25034417>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: ANP32B Antibody [NB100-1199] - Effect of overexpression or knockdown of PAL31 in C6 on H₂O₂-induced toxicity. (A) Representative micrographs showing C6 overexpressing GFP or GFP-tagged PAL31 after being treated with H₂O₂ (0 ~ 1 mM) for 4 hours. (B) MTT assay in GFP- or GFP-PAL31-overexpressed C6 after H₂O₂ treatment showing significant difference (cytoprotective effect of PAL31) at 0.5 mM & 1 mM H₂O₂ treatment between C6/GFP & C6/PAL31 groups. The data in each dosage were analyzed by two-way ANOVA & Bonferroni post hoc test. *P < 0.05, GFP (+H₂O₂) compared to PAL31 (+H₂O₂), n = 4, at 0.5 mM & 1 mM. (C) Western blot analysis of pal31 siRNA-treated C6 showing knockdown of PAL31 expression by PAL31 siRNA using 41.5 (lane1), 83 (lane2), & 166 (lane 3) picomole of pal31 siRNA or 332 picomole negative control. Actin works as a loading control. (D) MTT assay in Negative- or PAL31 silencer transfected C6 after H₂O₂ treatment showing significant difference at 1 mM H₂O₂ treatment between C6/Negative & C6/PAL31 siRNA groups. *P < 0.05, Negative (+H₂O₂) compared to PAL31 siRNA (+H₂O₂), n = 4. Magnification 100X (A). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/25034417>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: ANP32B Antibody [NB100-1199] - Overexpression of PAL31 in mixed glial culture effectively reduced LPS/IFN γ stimulation after co-culture with spinal cord neuron-glia cultures. (A) ED-1- immunoreactive microglia/macrophage in confluent mixed glial cultures (B) S100- immunoreactive astroglia in confluent mixed glial cultures (C) LPS/IFN γ -stimulated neuron-glia co-cultures (Glia/GFP + neuron) showing expression of both iNOS & GFP (D) LPS/IFN γ -stimulated neuron-glia co-cultures (Glia/PAL31+ neuron) showing expression of both iNOS & PAL31. (E) Quantitative analysis of released nitric oxide, as a form of nitrite, in LPS/IFN γ -stimulated or non-stimulated neuron-glia co-culture. Mixed glial cultures were transfected with pGFP or pPAL31 before seeding spinal cord neurons. Co-cultures were then treated with LPS/IFN γ for 2 days. Medium were saved for nitrite level determination, while cells were processed for immunostaining with iNOS. *P < 0.05, n = 4. PAL31 overexpressing co-cultures (+LPS/IFN γ) compared to GFP overexpressing co-cultures (+LPS/IFN γ). Magnification 200X (A-D). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/25034417>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: ANP32B Antibody [NB100-1199] - Overexpression of PAL31 in C6 did not affect the cell proliferative activity. (A) GFP/bright field-superimposed micrograph in GFP overexpressing cells showing > 50% transfective efficiency (B) GFP-PAL31/bright field-micrograph in PAL31 overexpressing C6 showing correct nuclear localization of GFP-tagged PAL31 & transfection efficiency. (C) Western blot analysis showing the expression of GFP (~27 kDa), GFP-tagged PAL31 (~58 kDa), PCNA (~34 kDa), endogenous PAL31 (~31 kDa) & Actin (~43 kDa) in GFP- or PAL31-C6 cultures. The level of PCNA, a protein expressing in nuclear during DNA synthesis, in cells did not alter after overexpression of GFP or GFP-tagged PAL31. Actin works as a loading control. (D) MTT assay in the transfected cells at 4 different time intervals (from Ctrl to 72 hours). Ctrl represents the cells after subculture & overnight incubation. The data in each time points were analyzed by two-way ANOVA & Bonferroni post hoc test. No significance, compared GFP & GFP tagged PAL31 groups. n = 3. Magnification 100X (A-B). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/25034417>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Zhu L, Perlaky L, Henning D, Valdez BC. Cloning and characterization of a new silver-stainable protein SSP29, a member of the LRR family. *Biochem Mol Biol Int* 1997-08-01 [PMID: 9285060]

Tseng Fan-Wei, Liou Dann-Ying, Tsai May-Jywan et al. Cytoprotective and anti-inflammatory effects of PAL31 overexpression in glial cells. *J Biomed Sci*. 2014-07-17 [PMID: 25034417] (WB, Rat)

NOVUS
BIOLOGICALS®
a biotechne® brand



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112

USA

Phone: 303.730.1950

Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400

Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane

Abingdon Science Park

Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15

Fax: (44) (0) 1235 533420

info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-1199

NB820-59242	Human Lymph Node Whole Tissue Lysate (Adult Whole Normal)
NB100-1199PEP	ANP32B Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1199

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

