

# Product Datasheet

## Fc gamma RII/CD32 Antibody NB100-1532

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-1532](http://www.novusbio.com/NB100-1532)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-1532](http://www.novusbio.com/reviews/destination/NB100-1532)



**NB100-1532**

## Fc gamma RII/CD32 Antibody

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Description	Novus Biologicals Goat Fc gamma RII/CD32 Antibody (NB100-1532) is a polyclonal antibody validated for use in IHC, WB, ELISA and ICC/IF. Anti-Fc gamma RII/CD32 Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	2213
Gene Symbol	FCGR2B
Species	Human
Specificity/Sensitivity	This antibody is expected to recognise all reported isoforms (NP_001002273.1; NP_001002274.1; NP_001002275.1; NP_003992.3; NP_001177757.1).
Immunogen	Peptide with sequence C-PDALEEPDDQNRI, from the C Terminus according to NP_003992.3; NP_001002273.1; NP_001002274.1; NP_001002275.1; NP_001177757.1.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 0.3-1 ug/ml, ELISA 1:32000, Immunohistochemistry 6 - 8 ug/ml, Immunocytochemistry/ Immunofluorescence 10 ug/ml, Immunohistochemistry-Paraffin 6 - 8 ug/ml
Application Notes	WB: Approx 38 kDa band observed in Daudi lysates (calculated MW of 34.0 kDa according to NP_003992.3).

## Images


Western Blot: Fc gamma RII/CD32 Antibody [NB100-1532] - (1ug/ml) staining of Daudi cell lysate (RIPA buffer, (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

250kDa  
150kDa  
100kDa  
75kDa  
50kDa  
37kDa  
25kDa  
20kDa  
15kDa


Immunohistochemistry-Paraffin: Fc gamma RII/CD32 Antibody [NB100-1532] - Staining (8ug/ml) of paraffin embedded Human Placenta. Heat induced antigen retrieval with citrate buffer pH 6, HRP-Staining.



Immunohistochemistry-Paraffin: Fc gamma RII/CD32 Antibody [NB100-1532] - Negative Control showing staining of paraffin embedded Human Placenta, with no primary antibody.



Immunocytochemistry/Immunofluorescence: Goat Polyclonal Fc gamma RII/CD32 Antibody [NB100-1532] - Immunofluorescence analysis of paraformaldehyde fixed THP-1 cells immobilized on Shifix™ coverslip, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



## Publications

Salas TR, Petrusseva I, Lavrik O et al. Evidence for direct contact between the RPA3 subunit of the human replication protein A single-strand DNA. *Nucleic Acids Res* 37(1):38-46. 2009-01-01 [PMID: 19010961]

McGaha TL, Sorrentino B, Ravetch JV. Restoration of tolerance in lupus by targeted inhibitory receptor expression. *Science*;307(5709):590-3. 2005-01-28 [PMID: 15681388]





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-1532**

---

NBL1-10656	Fc gamma RIIB/CD32b Overexpression Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-1532](http://www.novusbio.com/reviews/submit/NB100-1532)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)



