

Product Datasheet

Matrin 3 Antibody NB100-1761

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1 **Publications: 13**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1761

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1761



NB100-1761

Matrin 3 Antibody


Product Information	
Unit Size	0.1 ml
Concentration	0.2 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	TBS and 0.1% BSA

Product Description	
Description	Novus Biologicals Knockout (KO) Validated Rabbit Matrin 3 Antibody (NB100-1761) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-Matrin 3 Antibody: Cited in 13 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	9782
Gene Symbol	MATR3
Species	Human, Mouse
Immunogen	The immunogen recognized by this antibody maps to a region between residue 800 and the C-terminus (residue 847) of human Matrin 3 using the numbering given in entry NP_061322.2 (GeneID 9782).


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Knockout Validated
Recommended Dilutions	Western Blot 1:5000-1:15000, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence 1:250-1:1000, Immunoprecipitation 1-4 ug/mg of lysate, Immunohistochemistry-Paraffin 1:10-1:500, Knockout Validated
Application Notes	Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.

Images


Western blot shows lysates of HAP1 human near-haploid cell line and Matrin 3 knockout HAP1 cell line (KO). Nitrocellulose membrane was probed with Matrin 3 Antibody (Catalog # NB100-1761) O/N at 4C, followed by HRP-conjugated Secondary Antibody and ECL detection. A specific band was detected for Matrin 3 (as indicated) in the parental HAP1 cell line, but is not detectable in knockout HAP1 cell line. Primary antibody dilution used: 1/5000. The Ponceau stained transfers of each blot are shown. Image, protocol and testing courtesy of YCharOS Inc. (ycharos.com).




HAP1 lysates were prepared, and immunoprecipitation was performed using 2.0 ug of Matrin 3 antibody (Catalog # NBP1-04340) pre-coupled to Dynabeads protein A. Samples were washed and processed for Western Blot with Matrin 3 antibody. For Western Blot, Rb x Matrin 3 was used at 1/500. The Ponceau stained transfers of each blot are shown. SM=4% starting material; UB=4% unbound fraction; IP=immunoprecipitate. Image, protocol and testing courtesy of YCharOS Inc. (ycharos.com).




Immunoprecipitation: Matrin 3 Antibody [NB100-1761] - Detection of Human and Mouse Matrin 3 on HeLa whole cell lysate using NB100-1761. Matrin 3 was also immunoprecipitated using another rabbit anti-Matrin 3 antibody and NB100-1559.




Immunohistochemistry: Matrin 3 Antibody [NB100-1761] - Sample: FFPE section of human stomach carcinoma. Antibody: Affinity purified rabbit anti-Matrin 3 used at a dilution of 1:400 (0.5ug/ml). Detection: Red-fluorescent Goat anti-Rabbit IgG-heavy and light chain cross-adsorbed Antibody DyLight 594 Conjugated




Immunohistochemistry-Paraffin: Matrin 3 Antibody [NB100-1761] - Section of human laryngeal squamous cell carcinoma (left) and mouse squamous cell carcinoma (right). Antibody: Affinity purified rabbit anti-Matrin 3 used at a dilution of 1:1,000 (0.2ug/ml). Detection: DAB




Western Blot: Matrin 3 Antibody [NB100-1761] - The identification of host proteins immunoprecipitated with L1 ORF1p-FLAG. Validation of putative ORF1p-FLAG interacting proteins: Western blot images of the pJM101/L1.3FLAG and pJM101/L1.3 immunoprecipitation (IP) reactions. The pCEP4 lanes denote whole cell lysates derived from HeLa cells transfected with an empty pCEP4 vector (~ 1.0% input). Primary antibodies used to probe western blots are indicated to the left of the images. Immunoprecipitation reactions were conducted in either the absence (left) or presence (right) of RNaseA (10 ug/mL). The putative cellular functions of the ORF1p-FLAG interacting proteins are indicated on the right hand side of the blots. Image collected and cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pgen.1005121>) licensed under a CC-BY license.




Immunohistochemistry: Matrin 3 Antibody [NB100-1761] - Striking muscle pathology of gastrocnemius in Tg mice from Fo. 1579. Matrin 3 immunohistochemistry of transverse and longitudinal sections. * indicates rounded fiber. Arrow head indicates subsarcolemmal nucleus. Arrow indicates nuclear chain. Scale bars measure 25 um Image collected and cropped by CiteAb from the following publication (<https://actaneurocomms.biomedcentral.com/articles/10.1186/s40478-016-0393-5>), licensed under a CC-BY license.




Immunohistochemistry: Matrin 3 Antibody [NB100-1761] - Striking muscle pathology of gastrocnemius in Tg mice from Fo. 1579. Matrin 3 immunohistochemistry of transverse sections from phenotypic Tg show an increase in Matrin 3 immunoreactivity compared to NT transverse and longitudinal sections. * indicates rounded fiber. Arrow head indicates subsarcolemmal nucleus. Arrow indicates nuclear chain. Scale bars measure 25 um Image collected and cropped by CiteAb from the following publication (<https://actaneurocomms.biomedcentral.com/articles/10.1186/s40478-016-0393-5>), licensed under a CC-BY license.




Immunohistochemistry: Matrin 3 Antibody [NB100-1761] - Striking muscle pathology of gastrocnemius in Tg mice from Fo. 1579. Matrin 3 immunohistochemistry of f longitudinal sections from phenotypic Tg show an increase in Matrin 3 immunoreactivity compared to NT transverse and longitudinal sections. * indicates rounded fiber. Arrow head indicates subsarcolemmal nucleus. Arrow indicates nuclear chain. Scale bars measure 25 um Image collected and cropped by CiteAb from the following publication (<https://actaneurocomms.biomedcentral.com/articles/10.1186/s40478-016-0393-5>), licensed under a CC-BY license.




Immunohistochemistry: Matrin 3 Antibody [NB100-1761] - Striking muscle pathology of gastrocnemius in Tg mice from Fo. 1579. Matrin 3 immunohistochemistry of NT transverse and longitudinal sections. * indicates rounded fiber. Arrow head indicates subsarcolemmal nucleus. Arrow indicates nuclear chain. Scale bars measure 25 um Image collected and cropped by CiteAb from the following publication (<https://actaneurocomms.biomedcentral.com/articles/10.1186/s40478-016-0393-5>), licensed under a CC-BY license.




Immunoprecipitation: Matrin 3 Antibody [NB100-1761] - IP of human fibroblasts. Image from verified customer review.



HAP1 WT and Matrin 3 KO cells were labelled with a green or a far-red fluorescent dye, respectively. Cells were stained with Rab1A antibody and with the corresponding Alexa-fluor 555 coupled secondary antibody including DAPI. Acquisition of the blue (nucleus-DAPI), green (identification of WT cells), red (antibody staining) and far-red (identification of KO cells) channels was performed. Representative images of the blue and red (grayscale) channels are shown. WT and KO cells are outlined with green and magenta dashed line, respectively. Antibody dilution used: 1/200. Image, protocol and testing courtesy of YCharOS Inc. (ycharos.com).



Western Blot: Matrin 3 Antibody [NB100-1761] - TRAP150 inhibits the RRM-dependent binding activity of PSF, & inhibition of binding requires more than the minimal RRM binding domain. (A) UV crosslinking of ESS1 RNA with full-length PSF (PSF-FL) either alone (-), in the presence of BSA as a control or in the presence of increasing amounts of full-length FLAG-tagged TRAP150. (B) Same as panel (A), but competing binding of exRRMs of PSF with indicated truncations of GST-TRAP150 or GST alone. (C) Same as panel (A) but competing for binding of the minRRMs & hnRNP L by GST-TRAP(PID). (D) Western blots of immunoprecipitation of PSF from unstimulated (TRAP150 bound) & stimulated (TRAP150 unbound) JSL1 cells showing relative binding of TRAP150 & other known PSF-interacting partners. The source of the doublet for PSPC1 & p54nrb/NONO in stimulated cells is unknown, but is the same in input & IP samples. Image collected & cropped by CiteAb from the following publication (<https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkv816>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Guo Q, Han S The effect on m6A methylation writer complex by the reduced MATR3 in pterygium *Acta histochemica* 2023-10-30 [PMID: 37913560]

Haikerwal A, Barrera MD, Bhalla N et al. Inhibition of Venezuelan Equine Encephalitis Virus Using Small Interfering RNAs *Viruses* 2022-07-26 [PMID: 35893693] (IP, KD, WB, Human)

Details:

See Supplementary figure for KD

Moloney C, Rayaprolu S, Howard J et al. Analysis of spinal and muscle pathology in transgenic mice overexpressing wild-type and ALS-linked mutant MATR3 *Acta Neuropathol Commun* 2018-12-19 [PMID: 30563574] (WB, Mouse)

Marangi G, Lattante S, Doronzio PN et al. Matr3 variants are frequent in Italian ALS patients. *Neurobiol Aging* 2017-01-01 [PMID: 28029397]

Rayaprolu S, D'Alton S, Crosby K et al. Heterogeneity of Matr3 in the developing and aging murine central nervous system. *J Comp Neurol* 2016-10-01 [PMID: 26878116] (Mouse)

Moloney C, Rayaprolu S, Howard J et al. Transgenic mice overexpressing the ALS-linked protein Matr3 develop a profound muscle phenotype. *Acta Neuropathol Commun*. 2016-11-18 [PMID: 27863507] (WB, Mouse)

Christopher A. Yarosh, Iulia Tapescu, Matthew G. Thompson et al. TRAP150 interacts with the RNA-binding domain of PSF and antagonizes splicing of numerous PSF-target genes in T cells. *Nucleic Acid Research* 2015-08-10 [PMID: 26261210] (WB, Human)

Depreux FF, Puckelwartz MJ, Augustynowicz A et al. Disruption of the lamin A and matr3 interaction by myopathic LMNA mutations. *Hum. Mol. Genet.* 2015-05-06 [PMID: 25948554] (IP, WB, Mouse)

Moldovan JB, Moran JV. The Zinc-Finger Antiviral Protein ZAP Inhibits LINE and Alu Retrotransposition. *PLoS Genet* 2015-05-01 [PMID: 25951186] (ICC/IF, WB, Human)

Lee YW, Terranova C, Birkaya B et al. A novel nuclear FGF Receptor-1 partnership with retinoid and Nur receptors during developmental gene programming of embryonic stem cells. *J Cell Biochem* 2012-09-01 [PMID: 22539306]

Salton M, Elkon R, Borodina T et al. Matr3 binds and stabilizes mRNA. *PLoS One* 2011-01-01 [PMID: 21858232]

Erazo A, Yee MB, Banfield BW et al. The alpha herpesvirus US3/ORF66 protein kinases direct phosphorylation of the nuclear matrix protein matr3. *J Virol* 2011-01-01 [PMID: 20962082]

More publications at <http://www.novusbio.com/NB100-1761>



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1761

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1761

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

