

Product Datasheet

MIF Antibody NB100-1789

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1789

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1789



NB100-1789

MIF Antibody

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Description	Novus Biologicals Goat MIF Antibody (NB100-1789) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-MIF Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	4282
Gene Symbol	MIF
Species	Human, Rat
Reactivity Notes	Rat reactivity reported in scientific literature (PMID: 19066630).
Immunogen	Peptide with sequence C-NAANVGWNNSTFA corresponding to C-Terminus according to NP_002406.1.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA
Recommended Dilutions	Western Blot 0.01 - 0.03 ug/mL, Immunohistochemistry, Immunohistochemistry-Paraffin 5 ug/mL, Peptide ELISA 1:128000

Images


Western Blot: MIF Antibody [NB100-1789] - Staining of Human Thymus lysate (35 ug protein in RIPA buffer). Antibody at 0.01 ug/mL. Detected by chemiluminescence.




Immunohistochemistry: MIF Antibody [NB100-1789] - Co-localization of CXCR4 and MIF in urothelium. Sections from rats treated with saline or CYP are shown. MIF immunostaining (green immunofluorescence), CXCR4 immunostaining (red immunofluorescence) and an overlay panel combining both immunostaining and a DAPI nuclear stain. MIF immunostaining is seen in basal and intermediate cells and in fibroblasts in the lamina propria of saline treated rats, while superficial cells do not stain for MIF. Arrows show luminal edge of urothelium. CXCR4 is restricted to basal and intermediate cells of urothelium. CYP treatment resulted in superficial cell staining for MIF and CXCR4 and overlay panels demonstrate co-localization as orange color in urothelial cells. Arrows point to superficial cells showing MIF-CXCR4 co-localization. Calibration bar=50 μ m. Image collected and cropped by CiteAb from the following publication ([//dx.plos.org/10.1371/journal.pone.0003898](https://doi.org/10.1371/journal.pone.0003898)) licensed under a CC-BY license.




Western Blot: MIF Antibody [NB100-1789] - Staining of Daudi (A) and Jurkat (B) lysate (35 μ g protein in RIPA buffer). Antibody at 0.01 μ g/mL. Detected by chemiluminescence.



Immunohistochemistry-Paraffin: MIF Antibody [NB100-1789] - Staining of paraffin embedded Human Thymus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining. Antibody at 5 μ g/mL.



Immunocytochemistry/ Immunofluorescence: MIF Antibody [NB100-1789] - Co-localization of CXCR4 & MIF in urothelium. Representative sections from rats treated with saline (A–C) or CYP (D–I) are shown. The figure shows MIF immunostaining (green immunofluorescence), CXCR4 immunostaining (red immunofluorescence) & an overlay panel combining both immunostaining & a DAPI nuclear stain. MIF immunostaining is seen in basal & intermediate cells & in fibroblasts in the lamina propria of saline treated rats (A), while superficial cells do not stain for MIF. Arrows show luminal edge of urothelium. CXCR4 is restricted to basal & intermediate cells of urothelium (B) & lamina propria is not stained. Overlay of these panels (C) demonstrate co-localization of MIF & CXCR4 as orange coloring of cells. CYP treatment resulted in superficial cell staining for MIF (D,G) & CXCR4 (E,H) & overlay panels (F,I) demonstrate co-localization as orange color in urothelial cells. Arrows point to superficial cells showing MIF-CXCR4 co-localization. Calibration bar=50 μ m. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/19066630>), licensed under a CC0-1.0 license. Not internally tested by Novus Biologicals.



Publications

Church TS, Willis MS, Priest EL et al. Obesity, macrophage migration inhibitory factor, and weight loss. *Int J Obes Relat Metab Disord* 2005-03-29 [PMID: 15795748]

Vera PL, Iczkowski KA, Wang X, Meyer-Siegler KL. Cyclophosphamide-induced cystitis increases bladder CXCR4 expression and CXCR4-macrophage migration inhibitory factor association. *PLoS One* 2008-01-01 [PMID: 19066630] (Rat)

Hsieh, Y et al. CELL CYCLE, CELL DEATH, SENESENCE: Hepatitis B Virus Pre-S2 Mutant Surface Antigen Induces Degradation of Cyclin-Dependent Kinase Inhibitor p27Kip1 through c-Jun Activation Domain-Binding Protein 1. *Mol Cancer Res* 6:1063-1072. [PMID: 17951406]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1789

NB820-59242	Human Lymph Node Whole Tissue Lysate (Adult Whole Normal)
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1789

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



