

Product Datasheet

Pyruvate Dehydrogenase Kinase 1/PDK1 Antibody - BSA Free NB100-2384

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-2384

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-2384



NB100-2384**Pyruvate Dehydrogenase Kinase 1/PDK1 Antibody - BSA Free**


Product Information	
Unit Size	0.1 ml
Concentration	0.5 mg/ml
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Glycine and 0.15M NaCl
Target Molecular Weight	49 kDa

Product Description	
Description	Novus Biologicals Rabbit Pyruvate Dehydrogenase Kinase 1/PDK1 Antibody - BSA Free (NB100-2384) is a polyclonal antibody validated for use in WB. Anti-Pyruvate Dehydrogenase Kinase 1/PDK1 Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	5163
Gene Symbol	PDK1
Species	Human, Mouse
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 25925251).
Immunogen	A synthetic peptide made to an internal portion of the human PDK1 protein sequence (between residues 300-400). [Swiss-Prot Q15118]

Product Application Details	
Applications	Western Blot, Immunoblotting
Recommended Dilutions	Western Blot 0.5-2 ug/ml, Immunoblotting reported in scientific literature (PMID 28366875)
Application Notes	This PDK1 antibody is useful for Western blot, where a band is seen at ~49 kDa representing PDK1 protein. There are also two higher and one lower non-specific molecular weight bands. The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.

Images

Western Blot: PDK-1 Antibody [NB100-2384] - Detection of PDK-1 in human heart lysate using NB 100-2384 at 0.5 ug/ml.



Publications

Pastor-Soler NM, Sutton TA, Mang HE et al. Muc1 is protective during kidney ischemia-reperfusion injury Am J Physiol Renal Physiol. 2015-04-28 [PMID: 25925251] (WB, Mouse)

Gibier JB, Hemon B, Fanchon M et al. Dual role of MUC1 mucin in kidney ischemia-reperfusion injury: Nephroprotector in early phase, but pro-fibrotic in late phase. Biochim. Biophys. Acta. 2017-03-31 [PMID: 28366875] (IB)

Ho N, Coomber BL. Pyruvate dehydrogenase kinase expression and metabolic changes following dichloroacetate exposure in anoxic human colorectal cancer cells. Exp Cell Res 2015-02-01 [PMID: 25536473] (Human)

Lin HP, Jiang SS, Chuu CP. Caffeic Acid Phenethyl Ester Causes p21 Induction, Akt Signaling Reduction, and Growth Inhibition in PC-3 Human Prostate Cancer Cells. PLoS One. 2012-02-07 [PMID: 22347457]



Procedures

Western Blot Protocol for PDK1 Antibody (NB100-2384)

Pyruvate Dehydrogenase Kinase 1/PDK1 Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12%) on samples to be analyzed, loading 40 ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Rinse membrane with dH₂O and then stain the blot using ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% non-fat dry milk + 1% BSA in TBS for 1 hour at room temperature (RT).
6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
7. Dilute the rabbit anti-PDK1 primary antibody (NB 100-2384) in blocking buffer and incubate 1 hour at RT.
8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturer's instructions (we used BioFX Super Plus ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-2384

NB820-59217	Human Heart Whole Tissue Lysate (Adult Whole Normal)
NB100-2384PEP	Pyruvate Dehydrogenase Kinase 1/PDK1 Antibody Blocking Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-2384

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



