

Product Datasheet

AlphaB Crystallin/CRYAB Antibody - BSA Free NB100-2519

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-2519

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-2519



NB100-2519**AlphaB Crystallin/CRYAB Antibody - BSA Free**


Product Information	
Unit Size	0.1 ml
Concentration	1.08 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)

Product Description	
Description	Novus Biologicals Rabbit AlphaB Crystallin/CRYAB Antibody - BSA Free (NB100-2519) is a polyclonal antibody validated for use in WB and ICC/IF. Anti-AlphaB Crystallin/CRYAB Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1410
Gene Symbol	CRYAB
Species	Human, Mouse
Reactivity Notes	Predicted to react with rat, sheep, rabbit, monkey, guinea pig, chicken, and pig based on 100% sequence homology. Immunogen sequence has 81% homology with Zebrafish.
Immunogen	A synthetic peptide made to an internal region of the human Crystallin AB protein (within residues 70-140). [Swiss-Prot P02511]


Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 0.5-1.0 ug/ml, Immunocytochemistry/ Immunofluorescence 1:50 - 1:150. Use reported in scientific literature
Application Notes	In Western blot analysis, a band is seen at ~20 kDa. There is also a band of greater intensity at ~40 kDa that may be an aggregated heteromeric complex. **CryAB aggregates with homologous proteins, including CryAA and the small heat shock protein HSP28, to form large heteromeric complexes.

Images


Western Blot: AlphaB Crystallin/CRYAB Antibody [NB100-2519] - Ethanol induces the expression of heat shock proteins (HSPs) in primary astrocyte culture. Increase in alphabeta-crystallin protein levels after treatment for 2 h with 60 mmol/L ethanol (EtOH) or 42C heat shock (HS) in primary astrocyte culture. alpha-tubulin used as an internal standard. The bar graph to the right of the immunoblots represents the quantification of immunoreactive bands intensities normalized to the internal standard, expressed in arbitrary optical density (OD) units. The data are the mean +/- SEM of normalized relative OD values analyzed by one-way ANOVA with Dunnett's multiple comparison post hoc test, n = 3 (significantly differently at the level of *P < 0.05, **P < 0.01, ***P < 0.001). Image collected and cropped by CiteAb from the following publication (<https://doi.wiley.com/10.1002/brb3.125>) licensed under a CC-BY license.



Immunocytochemistry/Immunofluorescence: AlphaB Crystallin/CRYAB Antibody [NB100-2519] - CRYAB antibody was tested in SHSH5Y cells with DyLight488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and Dylight 550 (red).



Western Blot: AlphaB Crystallin/CRYAB Antibody [NB100-2519] - Detection of CryAB using NB 100-2519 (0.5ug/ml). Lane 1: Human skeletal muscle Lane 2: Mouse skeletal muscle.



Publications

Shaw A, TOth B, Arianti R Et al. BMP7 increases UCP1-dependent and independent thermogenesis with a unique gene expression program in human neck area derived adipocytes *Pharmaceuticals* 2021-09-28 [PMID: 34832860]

Chen AT, Xiao Y, Tang X et al. Cross-platform analysis reveals cellular and molecular landscape of glioblastoma invasion *Neuro-oncology* 2022-07-28 [PMID: 35901838]

Pignataro L, Varodayan FP, Tannenholz LE et al. Brief alcohol exposure alters transcription in astrocytes via the heat shock pathway. *Brain Behav.* 2013-03-01 [PMID: 23533150] (ICC/IF, Mouse)

Liu L, Qi X, Chen Z et al. Targeting the IRE1alpha/XBP1 and ATF6 Arms of the Unfolded Protein Response Enhances VEGF Blockade to Prevent Retinal and Choroidal Neovascularization. *Am J Pathol* 2013-02-07 [PMID: 23395094]

Procedures

Protocol specific for Crystallin AB Antibody (NB100-2519)

AlphaB Crystallin/CRYAB Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12%, Bis-Tris) on samples to be analyzed, loading 40 ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Rinse membrane with dH₂O and then stain the blot using ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% non-fat dry milk + 1% BSA in TBS, overnight at 4 degrees Celsius.
6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
7. Dilute the rabbit anti-crystallin AB primary antibody (NB 100-2519) in blocking buffer and incubate 1 hour at room temperature.
8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturer's instructions (Pierce's ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-2519

NB820-59253	Human Skeletal Muscle Whole Tissue Lysate (Adult Whole Normal)
NB100-2519PEP	AlphaB Crystallin/CRYAB Antibody Blocking Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-2519

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



