

Product Datasheet

CD14 Antibody

NB100-2807

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-2807

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-2807



NB100-2807

CD14 Antibody

Product Information

Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description


Description	Novus Biologicals Goat CD14 Antibody (NB100-2807) is a polyclonal antibody validated for use in IHC, WB, ELISA, Flow and ICC/IF. Anti-CD14 Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	929
Gene Symbol	CD14
Species	Human
Immunogen	Peptide with sequence C-KRVDADADPRQYAD corresponding to internal region according to NP_000582.1.

Product Application Details


Applications	Western Blot, ELISA, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1 - 3 ug/ml, Flow Cytometry 10 ug/ml, ELISA Detection limit 1:128000, Immunohistochemistry 6-8 ug/ml, Immunocytochemistry/ Immunofluorescence
Application Notes	WB: Approx 55kDa band observed in Human Lymph node and Tonsil lysates (calculated MW of 40.1kDa according to NP_000582.1). The observed molecular weight corresponds to the glycosylated form. Primary incubation 1 hour at room temperature. ICC/IF reactivity reported in (PMID: 25780044).

Images


Western Blot: CD14 Antibody [NB100-2807] - Staining (1ug/ml) of PBM lysate (35ug protein in RIPA buffer) with (B) and without (A) blocking with the immunizing peptide. Detected by chemiluminescence.



Flow Cytometry: CD14 Antibody [NB100-2807] - Flow cytometric analysis of paraformaldehyde fixed A549 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



Immunohistochemistry: Goat Polyclonal CD14 Antibody [NB100-2807] - (8 μ g/ml) staining of paraffin embedded Human Tonsil. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



Control: Goat Polyclonal CD14 Antibody [NB100-2807] - Negative Control showing staining of paraffin embedded Human Tonsil, with no primary antibody.



Publications

Iwaki D, Nishitani C, Mitsuzawa H et al. The CD14 region spanning amino acids 57-64 is critical for interaction with the extracellular Toll-like receptor 2 domain. *Biochem Biophys Res Commun* 2005-03-04 [PMID: 15670766]

Martin I, Caban-Hernandez K, Figueroa-Santiago O, Espino AM. *Fasciola hepatica* Fatty Acid Binding Protein Inhibits TLR4 Activation and Suppresses the Inflammatory Cytokines Induced by Lipopolysaccharide In Vitro and In Vivo. *J. Immunol.* 2015-03-16 [PMID: 25780044] (ICC/IF, WB, Human)

Funda DP, Tuckova L, Farre MA et al. CD14 is expressed and released as soluble CD14 by human intestinal epithelial cells in vitro: lipopolysaccharide activation of epithelial cells revisited. *Infect Immun* 2001-06-01 [PMID: 11349042]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-2807

NB820-59242	Human Lymph Node Whole Tissue Lysate (Adult Whole Normal)
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-2807

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

www.novusbio.com



technical@novusbio.com